

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A01N 43/04, A61K 39/02		A1	(11) International Publication Number: WO 99/13720 (43) International Publication Date: 25 March 1999 (25.03.99)
(21) International Application Number: PCT/US98/19600 (22) International Filing Date: 18 September 1998 (18.09.98)		(81) Designated States: AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>	
(30) Priority Data: 60/059,353 19 September 1997 (19.09.97) US (71) Applicant: THE OHIO STATE UNIVERSITY [US/US]; 1960 Kenny Road, Columbus, OH 43210 (US). (72) Inventors: RIKIHISA, Yasuko; 1120 Woodman Drive, Worthington, OH 43085 (US). OHASHI, Noris; 1210 Chambers Road, Columbus, OH 43212 (US). (74) Agent: DOCHERTY, Pamela, A.; Calfee, Halter & Griswold LLP, 1400 McDonald Investment Center, 800 Superior Avenue, Cleveland, OH 44114 (US).			
<p>(54) Title: OUTER MEMBRANE PROTEIN OF EHRLICHIA CANIS AND EHRLICHIA CHAFFEENSIS</p> <p>(57) Abstract</p> <p>The present invention relates to diagnostic tools for veterinary and human use which are used for serodiagnosing ehrlichiosis in mammals, particularly in members of the Canidae family and in humans. The present invention also provides polynucleotides which encode the outer membrane proteins of <i>E. chaffeensis</i>. The polynucleotides encode an OMP-1 family of proteins of <i>E. chaffeensis</i> and P30 family of proteins of <i>E. canis</i>. The present invention also provides the following isolated proteins of <i>E. chaffeensis</i> OMP-1, OMP-1A, OMP-1B, OMP-1C, OMP-1D, OMP-1E, OMP-1F, OMP-1R, OMP-1S, OMP-1T, OMP-1U, OMP-1V, OMP-1W, OMP-1X, and OMP-1Z, referred to hereinafter collectively as the "OMP family". The present invention also provides the following isolated proteins of <i>E. canis</i> P30, P30-a, P30-1, P30-2, P30-3, P30-4, P30-5, P30-6, P30-7, P30-8, P30-9, and P30-10, referred to hereinafter as the P30 family. The present invention also relates to an assay for diagnosing ehrlichiosis in humans using a recombinant outer membrane protein of <i>E. chaffeensis</i>, particularly OMP-1. The present invention also relates to an assay for diagnosing ehrlichiosis in humans and members of the family Canidae using a recombinant outer membrane protein of <i>E. canis</i>, particularly P30.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
RJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

**OUTER MEMBRANE PROTEIN OF EHRLICHIA CANIS
AND EHRLICHIA CHAFFEENIS**

This work was supported by grant RO1 AI40934 from National Institutes of Health. The government has certain rights in this invention.

BACKGROUND OF THE INVENTION

The ehrlichiae are obligate intracellular bacteria that infect circulating leucocytes. *Ehrlichia chaffeensis* infects the monocytes and macrophages in humans and causes human monocytic ehrlichiosis. The clinical manifestations of ehrlichiosis in humans are nonspecific and similar to Rocky Mountain spotted fever. The clinical manifestations include fever, chills, headache myalgia or vomiting and weight loss. Most patients have a history of tick exposure.

Ehrlichia canis infects and causes ehrlichiosis in animals belonging to the family Canidae. Canine ehrlichiosis consists of an acute and a chronic phase. The acute phase is characterized by fever, serous nasal and ocular discharges, anorexia, depression, and loss of weight. The chronic phase is characterized by severe pancytopenia, epistaxis, hematuria, blood in feces in addition to more severe clinical signs of the acute disease. If treated early during the course of the disease, dogs respond well to doxycycline. However, chronically infected dogs do not respond well to the antibiotic. Therefore, early diagnosis is very important for treating canine ehrlichiosis.

The primary diagnostic test for diagnosing canine ehrlichiosis and human ehrlichiosis is the indirect fluorescent antibody (IFA) test. This test uses the etiologic agent *Ehrlichia canis* to diagnose canine ehrlichiosis. The IFA test uses *Ehrlichia chaffeensis* as antigen for diagnosing human ehrlichiosis. The IFA test has, however, serious limitations. The IFA test is subject to false positives because the antigens are made of whole infected cells which comprise many nonspecific proteins which will cross-react with sera from some patients. The IFA test is also subject to false negatives because IFA antigens are unstable and may become inactivated during storage. In addition the IFA test requires a special equipment to perform the test. For example, the IFA test requires a tissue culture system for growing the bacterium that are used to prepare the antigen slides, a fluorescent microscope, and trained persons to evaluate the serum reactivity to the bacterial antigen on the slide.

Tools which permit simpler, more rapid, and objective serodiagnosis of canine ehrlichiosis or human ehrlichiosis are desirable.

SUMMARY OF THE INVENTION

The present invention relates to improved diagnostic tools for veterinary and human use which are used for serodiagnosing ehrlichiosis in mammals, particularly in members of the Canidae family and in humans.

The present invention also provides polynucleotides or nucleic acids which encode the outer membrane proteins of *E. chaffeensis*. The OMP-1 polynucleotide encodes an OMP-1 protein of *E. chaffeensis* having a molecular weight of about 27.7 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG.3B, SEQ ID NO: _____. The OMP-1B polynucleotide encodes an OMP-1B protein of *E.*

chafeensis having a molecular weight of about 28.2 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 4B, SEQ ID NO: ___. The OMP-1C polynucleotide encodes an OMP-1C protein of *E. chafeensis* having a molecular weight of about 27.6 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 5B, SEQ ID NO: ___. The OMP-1D polynucleotide encodes an OMP-1D protein of *E. chafeensis* having a molecular weight of about 28.7 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 6B, SEQ ID NO: ___. The OMP-1E polynucleotide encodes an OMP-1E protein of *E. chafeensis* having a molecular weight of about 27.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 7B, SEQ ID NO: ___. The OMP-1F polynucleotide encodes an OMP-1F protein of *E. chafeensis* having a molecular weight of about 27.9 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 8B, SEQ ID NO: ___. The OMP-1A polynucleotide encodes an OMP-1A protein of *E. chafeensis* having a molecular weight of about 29.6 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 9B, SEQ ID NO: ___. The OMP-1R polynucleotide encodes an OMP-1R protein of *E. chafeensis* having a molecular weight of at least 23 kDa and comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 10B, SEQ ID NO: ___. The OMP-1S polynucleotide encodes an OMP-1S protein of *E. chafeensis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 11B, SEQ ID NO: ___. The OMP-1T polynucleotide encodes an OMP-1T protein of *E. chafeensis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 12B, SEQ ID NO: ___. The OMP-1U polynucleotide encodes an OMP-1U protein of *E. chafeensis* having a molecular weight of about 30.6 kDa and an amino acid sequence which is at least 85% homologous to amino acid sequence shown in FIG. 13B, SEQ ID NO: ___. The OMP-1V polynucleotide encodes an OMP-1V protein of *E. chafeensis* having a molecular weight of about 28.0 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 14B, SEQ ID NO: ___. The OMP-1W polynucleotide encodes an OMP-1W protein of *E. chafeensis* having a molecular weight of about 28.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 15B, SEQ ID NO: ___. The OMP-1X polynucleotide encodes an OMP-1S protein of *E. chafeensis* having a molecular weight of about 27.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 16B, SEQ ID NO: ___. The OMP-1Y polynucleotide encodes an OMP-1Y protein of *E. chafeensis* having a molecular weight about 28.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 17B, SEQ ID NO: ___. The OMP-1Z polynucleotide encodes an OMP-1Z protein of *E. chafeensis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 18B, SEQ ID NO: ____.

The outer membrane proteins from *E. chaffeensis*, particularly a recombinant form of OMP-1, are immunogenic and, thus are useful for preparing antibodies. Such antibodies are useful for immunolabeling isolates of *E. chaffeensis* and for detecting the presence of *E. chaffeensis* in body fluids, tissues, and particularly in monocytes and macrophages. The isolated outer membrane proteins, particularly OMP-1, are also useful for

detecting antibodies to *E. chaffeensis* in the blood of patients with clinical signs of ehrlichiosis. The isolated outer membrane protein, particularly OMP-1, are also useful immunogens for raising antibodies that are capable of reducing the level of infection in an immunized mammal that has been infected with *E. chaffeensis*. The isolated membrane proteins are also useful in a vaccine for protecting against infection with *E. chaffeensis*.

The present invention also relates to isolated polynucleotides which encode 30 kDa outer membrane proteins from *Ehrlichia canis*. The proteins are designated P30 and P30a. The proteins, particularly P30, are immunogenic and are, thus, useful for preparing antibodies that are useful for immunolabeling isolates of *E. canis*. The P30 protein is also useful for diagnosing canine ehrlichiosis in mammals, particularly in members of the family Canidae, most particularly in dogs and for diagnosing infections with *E. chaffeensis* in humans. The P30 protein is also a useful immunogen for raising antibodies that reduce the level of infection in an immunized mammal that has been infected with *E. canis*. The P30 protein is also useful in a vaccine for protecting animals against infection with *E. canis*.

The present invention also provides the following isolated proteins of *E. chaffeensis* OMP-1 (also known as p28), OMP-1A, OMP-1B, OMP-1C, OMP-1D, OMP-1E, OMP-1F, OMP-1R, OMP-1S, OMP-1T, OMP-1U, OMP-1V, OMP-1W, OMP-1X, and OMP-1Z, referred to hereinafter collectively as the "OMP family". The present invention also provides the following isolated proteins of *E. canis* P30, P30-a, P30-1, P30-2, P30-3, P30-4, P30-5, P30-6, P30-7, P30-8, P30-9, and P30-10, referred to hereinafter as the P30 family.

The present invention also relates to an assay for diagnosing ehrlichiosis in humans using a recombinant outer membrane protein of *E. chaffeensis*, particularly OMP-1. The present invention also relates to an assay for diagnosing ehrlichiosis in humans and members of the family Canidae using a recombinant outer membrane protein of *E. canis*, particularly P30.

Brief Description of the Figures

FIG. 1. shows the DNA sequence of and the amino acid sequence encoded by the *E. chaffeensis* (p28) gene cloned in pCRII/p28. The N-terminal amino acid sequence of native *omp-1* protein (P28) determined chemically is underlined. Five amino acid residues at the N terminus of P28 which were not included in the p28 gene, are indicated by boldface. Arrows indicate annealing positions of the primer pair designed for PCR.

FIG. 2. shows the restriction map of 6.3-kb genomic DNA including the *omp-1* gene copies in *E. chaffeensis*. The four DNA fragments were cloned from the genomic DNA (pPS2.6, pPS3.6, pEC2.6, and pEC3.6). A recombinant plasmid pPS2.6 has an overlapping sequence with that of pEC3.6. The closed boxes at the bottom show PCR-amplified fragments from the genomic DNA for confirmation of the overlapping area. Open boxes at the top indicate open reading frames (ORF) of *omp-1* gene copies with direction by arrows. Open boxes at the bottom show DNA fragments subcloned for DNA sequencing.

FIG. 3B shows one embodiment of the OMP-1 protein; FIG. 3A shows one embodiment of the OMP-1 polynucleotide.

FIG. 4B shows one embodiment of the OMP-1B protein, FIG. 4A shows one embodiment of the OMP-1B polynucleotide

FIG. 5A shows one embodiment of the OMP-1C p lynucleotide; FIG 5B shows one embodiment of the OMP-1C protein.

FIG. 6B shows one embodiment of the OMP-1D protein; FIG. 6A shows one embodiment of the OMP-1D polynucleotide.

FIG. 7A shows one embodiment of the OMP-1E protein; FIG 7B shows one embodiment of the OMP-1E polynucleotide.

FIG. 8A shows one embodiment of the OMP-1F protein; FIG 8 B shows one embodiment of the OMP-1F polynucleotide.

FIG. 9B shows one embodiment of the OMP-1A protein, FIG 9A shows one embodiment of the OMP-1A polynucleotide.

FIG. 10 B shows one embodiment of a portion of the OMP-1R protein, FIG 10A shows one embodiment of an OMP-1R polynucleotide encoding such polypeptide.

FIG. 11 B shows one embodiment of a portion of the OMP-1S protein, FIG 11A shows one embodiment of the OMP-1S polynucleotide encoding such polypeptide.

FIG. 12 B shows one embodiment of a portion of the OMP-1T protein, FIG 12A shows one embodiment of the OMP-1T polynucleotide encoding such polypeptide.

FIG. 13 B shows one embodiment of the OMP-1U protein, FIG 13A shows one embodiment of the OMP-1U polynucleotide.

FIG. 14 B shows one embodiment of the OMP-1V protein, FIG 14A shows one embodiment of the OMP-1V polynucleotide.

FIG. 15 B shows one embodiment of the OMP-1W protein, FIG 15A shows one embodiment of the OMP-1W polynucleotide.

FIG. 16 B shows one embodiment of the OMP-1X protein, FIG 16A shows one embodiment of the OMP-1X polynucleotide.

FIG. 17 B shows one embodiment of the OMP-1Y protein, FIG 17A shows one embodiment of the OMP-1Y polynucleotide.

FIG. 18 B shows one embodiment of the OMP-1Z protein, FIG 18A shows one embodiment of the OMP-1Z polynucleotide.

FIG. 19 B shows one embodiment of the P30 protein, FIG 19A shows one embodiment of the P30 polynucleotide.

FIG. 20 B shows one embodiment of the P30a protein, FIG 20A shows one embodiment of the p30A p lynucleotide.

FIG. 21 B shows one embodiment of the P30-1 protein, FIG 21A shows one embodiment of the p30-1 polynucleotide.

FIG. 22 B shows one embodiment of the P30-2 protein, FIG 22 A shows one embodiment of the p30-2 polynucleotide.

FIG. 23 B shows one embodiment of the P30-3 protein, FIG 23 A shows one embodiment of the p30-3 polynucleotide.

FIG. 24 B shows one embodiment of the P30-4 protein, FIG 22 A shows one embodiment of the p30-4 polynucleotide.

FIG. 25 B shows one embodiment of the P30-5 protein, FIG 22 A shows one embodiment of the p30-5 polynucleotide.

FIG. 26 B shows one embodiment of the P30-6 protein, FIG 26 A shows one embodiment of the p30-6 polynucleotide.

FIG. 27 B shows one embodiment of the P30-7 protein, FIG 27 A shows one embodiment of the p30-7 polynucleotide.

FIG. 28 B shows one embodiment of the P30-8 protein, FIG 28 A shows one embodiment of the p30-8 polynucleotide.

FIG. 29 B shows one embodiment of a portion of the P30-9 protein, FIG 29 A shows one embodiment of the p30-9 polynucleotide encoding such polypeptide.

FIG. 30 B shows one embodiment of a portion of the P30-10 protein, FIG 30 A shows one embodiment of the p30-10 polynucleotide encoding such polypeptide.

FIG. 31 depicts the amino acid sequences alignment of seven *E. chaffeensis* OMP-1s and *Cowdria ruminantium* MAP-1. Aligned positions of identical amino acids with OMP-IF are shown with dots. The sequence of *C. ruminantium* MAP-1 is from the report of Van Vliet et al (1994) Molecular cloning, sequence analysis, and expression of the gene encoding the immunodominant 32-kilodalton protein of *Cowdria ruminantium*. Infect. Immun. 62:1451-1456. Gaps indicated by dashes were introduced for optimal alignment of all proteins. Bars indicates semivariable region (SV) and three hypervariable regions (HV1, HV2, and HV3).

DETAILED DESCRIPTION OF THE INVENTION

Isolated Polynucleotides Encoding OMP-1, OMP-1A, OMP-1B, OMP-1C, OMP-1D, OMP-IF and the OMP from E. Canis

In one aspect, the present invention, provides isolated polynucleotides that encode the outer membrane proteins, OMP-1 (or p28), OMP-1B, OMP-1C, OMP-1D, OMP-1E, OMP-1F, OMP-1A, OMP-1R, OMP-1S, OMP-1T, OMP-1U, OMP-1V, OMP-1W, OMP-1X, OMP-1Y and OMP-1Z from *E. chaffeensis* and the outer membrane proteins P30, P30-a, P-30-1, P30-2, P30-3, P30-4, P30-5, P30-6, P30-7, P30-8, P30-9, and P30-10 from *E. Canis* or an immunogenic fragment thereof.

The polynucleotide is single stranded or double stranded. The polynucleotide may be a DNA or RNA molecule, preferably a DNA molecule, and comprises a sequence which codes for the respective outer membrane protein. Preferably, the polynucleotide encodes at least the mature form of outer membrane protein. The polynucleotide optionally further comprises a leader sequence and encode an outer membrane preprotein that is

processed in the cell to form the mature protein. The polynucleotide of the present invention may also be fused in frame to a marker sequence which allows for purification of the corresponding outer membrane protein.

The OMP-1 polynucleotide encodes an OMP-1 protein of *E. chafeensis* having a molecular weight of about 27.7 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 3B SEQ ID NO: ____; Figure 3B shows one embodiment of the OMP-1 protein. Figure 3A shows one embodiment of the OMP-1 polynucleotide. The OMP-1B polynucleotide encodes an OMP-1B protein of *E. chafeensis* having a molecular weight of about 28.2 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 4B SEQ ID NO: ____; Figure 4B shows one embodiment of the OMP-1B protein. Figure 4A shows one embodiment of the OMP-1B polynucleotide. The OMP-1C polynucleotide encodes an OMP-1C protein of *E. chafeensis* having a molecular weight of about 27.6 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 5B SEQ ID NO: ____; Figure 5B shows one embodiment of the OMP-1C protein. Figure 5A shows one embodiment of the OMP-1C polynucleotide. The OMP-1D polynucleotide encodes an OMP-1D protein of *E. chafeensis* having a molecular weight of about 28.7 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 6B SEQ ID NO: ____; Figure 6B shows one embodiment of the OMP-1D protein. Figure 6A shows one embodiment of the OMP-1D polynucleotide. The OMP-1E polynucleotide encodes an OMP-1E protein of *E. chafeensis* having a molecular weight of about 27.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 7B SEQ ID NO: ____; Figure 7B shows one embodiment of the OMP-1E protein. Figure 7A shows one embodiment of the OMP-1E polynucleotide. The OMP-1F polynucleotide encodes an OMP-1F protein of *E. chafeensis* having a molecular weight of about 27.9 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 8B SEQ ID NO: ____; Figure 8B shows one embodiment of the OMP-1F protein. Figure 8A shows one embodiment of the OMP-1F polynucleotide. The OMP-1A polynucleotide encodes an OMP-1A protein of *E. chafeensis* having a molecular weight of about 29.6 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 9B SEQ ID NO: ____; Figure 9B shows one embodiment of the OMP-1A protein. Figure 9A shows one embodiment of the OMP-1A polynucleotide. The OMP-1R polynucleotide encodes an OMP-1R protein of *E. chafeensis* having a molecular weight of at least 23 kDa and comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 10B SEQ ID NO: ____; Figure 10B shows one embodiment of a portion of the OMP-1R protein. Figure 10A shows one embodiment of the OMP-1R polynucleotide encoding such polynucleotide. The OMP-1S polynucleotide encodes an OMP-1S protein of *E. chafeensis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 11B SEQ ID NO: ____; Figure 11B shows one embodiment of a portion of the OMP-1S protein. Figure 11A shows one embodiment of the OMP-1S polynucleotide encoding such polypeptide. The OMP-1T polynucleotide encodes an OMP-1T protein of *E. chafeensis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 12B SEQ ID NO: ____; Figure 12B shows one embodiment of a portion of the OMP-1T protein. Figure 12A shows one embodiment of the OMP-1T polynucleotide encoding such polypeptide. The OMP-1U polynucleotide encodes an

OMP-1U protein of *E. chafeensis* having a molecular weight of about 30.6 kDa and an amino acid sequence which is at least 85% homologous to amino acid sequence shown in FIG. 13B SEQ ID NO: __; Figure 13B shows one embodiment of the OMP-1U protein, Figure 13A shows one embodiment of the OMP-1U polynucleotide. The OMP-1V polynucleotide encodes an OMP-1V protein of *E. chafeensis* having a molecular weight of about 28.0 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 14B SEQ ID NO: __; Figure 14B shows one embodiment of the OMP-1V protein, Figure 14A shows one embodiment of the OMP-1V polynucleotide. The OMP-1W polynucleotide encodes an OMP-1W protein of *E. chafeensis* having a molecular weight of about 28.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 15B SEQ ID NO: __; Figure 15B shows one embodiment of the OMP-1W protein, Figure 15A shows one embodiment of the OMP-1W polynucleotide. The OMP-1X polynucleotide encodes an OMP-1S protein of *E. chafeensis* having a molecular weight of about 27.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 16B SEQ ID NO: __; Figure 16B shows one embodiment of the OMP-1X protein, Figure 16A shows one embodiment of the OMP-1X polynucleotide. The OMP-1Y polynucleotide encodes an OMP-1Y protein of *E. chafeensis* having a molecular weight about 28.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 17B SEQ ID NO: __; Figure 17B shows one embodiment of the OMP-1Y protein, Figure 17A shows one embodiment of the OMP-1Y polynucleotide. The OMP-1Z polynucleotide encodes an OMP-1Z protein of *E. chafeensis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 18B SEQ ID NO: __; Figure 18B shows one embodiment of a portion of the OMP-1Z protein, Figure 18A shows one embodiment of an OMP-1Z polynucleotide encoding such polypeptide.

The p30 polynucleotide encodes a P30 protein of *E. canis* having a molecular weight of about 28.8 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 19B SEQ ID NO: __; Figure 19B shows one embodiment of the P30 protein, Figure 19A shows one embodiment of the p30 polynucleotide. The p30A polynucleotide encodes a P30a protein of *E. canis* having a molecular weight of about 29.1 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 20B SEQ ID NO: __; Figure 20B shows one embodiment of the P30a protein, Figure 20A shows one embodiment of the p30A polynucleotide. The p30-1 polynucleotide encodes a P30-1 protein of *E. canis* having a molecular weight of about 27.7 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 21B SEQ ID NO: __; Figure 21B shows one embodiment of the P30-1 protein, Figure 21A shows one embodiment of the p30-1 polynucleotide. The p30-2 polynucleotide encodes a P30-2 protein of *E. canis* having a molecular weight of about 28.0 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 22B SEQ ID NO: __; Figure 22B shows one embodiment of the P30-2 protein, Figure 22A shows one embodiment of the p30-2 polynucleotide. The p30-3 polynucleotide encodes a P30-3 protein of *E. canis* having a molecular weight of about 28.7 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 23B SEQ ID NO: __; Figure 23B shows one embodiment of the P30-3 protein, Figure 23A shows one embodiment of the p30-3 polynucleotide. The p30-4 polynucleotide

encodes a P30-4 protein of *E. canis* having a molecular weight of about 28.0 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 24B SEQ ID NO: ____; Figure 24B shows one embodiment of the P30-4 protein, Figure 24A shows one embodiment of the p30-4 polynucleotide. The p30-5 polynucleotide encodes a P30-5 protein of *E. canis* having a molecular weight of about 29.4 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 25B SEQ ID NO: ____; Figure 25B shows one embodiment of the P30-5a protein, Figure 25A shows one embodiment of the p30-5a polynucleotide. The p30-6 polynucleotide encodes a P30-6 protein of *E. canis* having a molecular weight of about 29.5 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 26B SEQ ID NO: ____; Figure 26B shows one embodiment of the P30-6 protein, Figure 26A shows one embodiment of the p30-6 polynucleotide. The p30-7 polynucleotide encodes a P30-7 protein of *E. canis* having a molecular weight of about 29.9 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 29B SEQ ID NO: ____; Figure 29B shows one embodiment of the P30-7 protein, Figure 29A shows one embodiment of the p30-7 polynucleotide. The p30-8 polynucleotide encodes a P30-8 protein of *E. canis* having a molecular weight of about 30.3 kDa and an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 28B SEQ ID NO: ____; Figure 28B shows one embodiment of the P30-8 protein, Figure 28A shows one embodiment of the p30-8 polynucleotide. The p30-9 polynucleotide encodes a P30-9 protein of *E. canis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 29B SEQ ID NO: ____; Figure 29B shows one embodiment of a portion of the P30-9 protein, Figure 29A shows one embodiment of the p30-9 polynucleotide encoding such polypeptide. The p30-10 polynucleotide encodes a P30-10 protein of *E. canis* comprising an amino acid sequence which is at least 85% homologous to the amino acid sequence shown in FIG. 30B SEQ ID NO: ____; Figure 30B shows one embodiment of a portion of the P30-10 protein, Figure 30A shows one embodiment of the p30-10 polynucleotide encoding such polypeptide.

The polynucleotides encoding an *E. chafeensis* outer membrane protein or an *E. canis* outer membrane protein have a sequence that is at least 85%, preferably at least 90%, more preferably at least 95% homologous to or similar to the amino acid sequences shown in Figures 3B through 30B, and thus embrace polynucleotides encoding outer membrane proteins from different strains of *E. chafeensis* and *E. canis*. The polynucleotides encode an outer membrane protein whose conserved regions collectively are at least 90%, preferably at 95%, more preferably at least 97% homologous to the conserved regions of the amino acid sequences of the present invention. The outer membrane proteins of *E. chafeensis* and *E. canis* have six conserved regions, which are separated by one semivariable region and three hypervariable regions. The conserved regions of the outer membrane proteins OMP-1, OMP-1A, OMP-1B, OMP1-C, OMP1-D, OMP1-F are depicted in Fig. 31. Preferably, the amino acid sequence of the outer membrane proteins of *E. chafeensis* and *E. canis* are at least 30% divergent from the amino acid sequence of MAP-1. Such sequences include allelic, strain variants and other amino acid sequence variants (e.g., including "muteins" or "mutant proteins"), whether naturally-occurring or biosynthetically produced. As used herein, "amino acid sequence homology" is understood to mean amino acid sequence similarity, and homologous sequences share identical or similar amino acids, where similar amino acids are conserved amino acids as defined by

Dayoff et al., *Atlas of Protein Sequence and Structure*; vol. 5, Supp. 3, pp. 345-362. (M. O. Dayoff, ed., Nat'l BioMed. Research Fdn., Washington D.C. 1978.) Thus, a candidate sequence sharing 85% amino acid sequence homology with a reference sequence requires that, following alignment of the candidate sequence with the reference sequence, 85% of the amino acids in the candidate sequence are identical to the corresponding amino acid in the reference sequence, or constitute a conserved amino acid change thereto. "Amino acid sequence identity" is understood to require identical amino acids between two aligned sequences. Thus, a candidate sequence sharing 85% amino acid identity with a reference sequence requires that, following alignment of the candidate sequence with the reference sequence, 85% of the amino acids in the candidate sequence are identical to the corresponding amino acid in the reference sequence.

As used herein, all homologies and identities are calculated using the amino acid sequences shown in the cited Figure or SEQ ID NO as the reference sequence. Thus, to determine whether an amino acid sequence is 85% homologous to OMP-1, one uses the amino acid sequence shown in Fig. ___, SEQ ID NO: __ as a reference.

Also as used herein, sequences are aligned for homology and identity calculations using the method of the software basic local alignment search tool in the BLAST network service (the National Center for Biotechnology Information, Bethesda, MD) which employs the method of Altschul, S. F., Gish, W., Miller, W., Myers, E. W. & Lipman, D. J. (1990) *J. Mol. Biol.* 215, 403-410. Identities are calculated by the Align program (DNAstar, Inc.) In all cases, internal gaps and amino acid insertions in the candidate sequence as aligned are ignored when making the homology/identity calculation.

In another aspect, the present invention provides a nucleotide sequence encoding a polypeptide which comprises a fragment of the OMP1 protein, hereinafter referred to as "rP28". The rP28 polypeptide weighs approximately 31 kDa and comprises all but of the first 5 amino acids of mature OMP-1 protein. The rP28 polypeptide comprises the amino acid sequence extending from amino acid 6 through amino acid 251 of the amino acid sequence shown in Fig. 1, SEQ ID NO. The present invention also embraces polypeptides where one or more of the amino acids in the sequence extending from amino acid 1 or 6 through amino acid 251 Fig. 1 are replaced by conservative amino acid residues. The present invention also relates to derivatives of rP28 that have an amino acid sequence identity of at least 85%, more preferably at least 90%, and most preferably of at least 95% with the amino acid sequence extending from amino acid 1 or 6 through amino acid 251 of the protein and which derivative binds to antibodies in sera from humans infected with *E. chaffeensis*.

The polynucleotides are useful for producing the outer membrane proteins of *E. chaffeensis* and *E. canis*. For example, an RNA molecule encoding the outer membrane protein OMP-1 is used in a cell-free translation systems to prepare OMP-1. Alternatively, a DNA molecule encoding the outer membrane protein is introduced into an expression vector and used to transform cells. Suitable expression vectors include for example chromosomal, nonchromosomal and synthetic DNA sequences, e.g., derivatives of SV40, bacterial plasmids, phage DNAs; yeast plasmids, vectors derived from combinations of plasmids and phage DNAs, viral DNA such as vaccinia, adenovirus, fowl pox virus, and pseudorabies. The DNA sequence is introduced into the expression vector by conventional procedures.

Accordingly, the present invention also relates to recombinant constructs comprising one or more of the polynucleotide sequences. Suitable constructs include, for example, vectors, such as a plasmid, phagemid, or viral vector, into which a sequence that encodes the outer membrane protein has been inserted. In the expression vector, the DNA sequence which encodes the outer membrane protein is operatively linked to an expression control sequence, i.e., a promoter, which directs mRNA synthesis. Representative examples of such promoters, include the LTR or SV40 promoter, the *E. coli* lac or trp, the phage lambda PL promoter and other promoters known to control expression of genes in prokaryotic or eukaryotic cells or in viruses. The promoter may also be the natural promoter of the outer membrane protein coding sequence. The expression vector also contains a ribosome binding site for translation initiation and a transcription terminator. Preferably, the recombinant expression vectors also include an origin of replication and a selectable marker, such as for example, the ampicillin resistance gene of *E. coli* to permit selection of transformed cells, i.e. cells that are expressing the heterologous DNA sequences. The polynucleotide sequence encoding the outer membrane protein is incorporated into the vector in frame with translation initiation and termination sequences. Optionally, the sequence encodes a fusion outer membrane protein which includes an N-terminal or C-terminal peptide or tag that stabilizes or simplifies purification of the expressed recombinant product. Representative examples of such tags include sequences which encode a series of histidine residues, the Herpes simplex glycoprotein D, or glutathione S-transferase.

Polynucleotides which encode portions of the outer membrane proteins of *E. chaffeensis* and *E. canis* are useful as probes for isolating and identifying *E. chaffeensis* genes and *E. canis* genes, particularly full-length genes from new strains or isolates of *E. chaffeensis* and *E. canis*.

The Outer Membrane Proteins of *E. chaffeensis* and *E. Canis*

In addition to the outer membrane proteins OMP-1, OMP-1B, OMP-1C, OMP-1D, OMP-1E, and OMP-1F, OMP-1R, OMP-1S, OMP-1T, OMP-1U, OMP-1V, OMP-1W, OMP-1X, OMP-1Y, and OMP-1Z from *E. chaffeensis* and the proteins P30, P30A, P30-1, P30-2, P30-3, P30-4, P30-5, P30-6, P30-7, P30-8, P30-9, and P30-10 from *E. Canis*, the present invention embraces non-naturally occurring allelic forms or derivatives of the outer membrane proteins, where one or more of the amino acids have been replaced by conservative amino acid residues, typically by using direct synthesis or recombinant techniques.

Preparing the Outer Membrane Proteins

The outer membrane proteins of the present invention are synthetically produced by conventional peptide synthesizers. The outer membrane proteins are also produced using cell-free translation systems and RNA molecules derived from DNA constructs that encode the outer membrane protein. Alternatively, the outer membrane protein is made by transfecting host cells with expression vectors that comprise a DNA sequence which encodes the outer membrane protein and then inducing expression of the outer membrane protein in the host cells.

The outer membrane protein is expressed in suitable host cells, preferably bacteria, under the control of suitable promoters. Host cells are transformed with the expression vectors of this invention and cultured in conventional nutrient media. Such media optionally contains additional compounds, such as for example

compounds that induce promoters, such as for example isopropyl- β -D-thi galactoside which induces the Lac promoter, or compounds, such as for example, ampicillin, which allows for selection of transformants.

Following transformation of the suitable host strain and growth of the host strain to an appropriate cell density, the cells are typically harvested by centrifugation, disrupted by physical or chemical means, and the resulting crude extract retained for further purification of the outer membrane protein. Such purification usually involves salting-out of the protein fraction, and one or more chromatography steps, including aqueous ion exchange chromatography, size exclusion chromatography steps, and high performance liquid chromatography (HPLC).

Preparation of Antibodies

The isolated outer membrane proteins, particularly the recombinant forms of the outer membrane proteins, are used as immunogens to produce antibodies immunospecific for the corresponding protein. The term "immunospecific" means the antibodies have substantially greater affinity for the protein used as an immunogen than for other proteins. Such antibodies are generated using conventional techniques by administering the respective outer membrane protein or a portion thereof, i.e., the recombinant polypeptide, to an animal, preferably a nonhuman, collecting blood from the immunized animals and isolating the serum and/or the IgG fraction from the blood. Monoclonal antibodies are prepared by injecting animals with the immunogens, extracting antibody-producing B cells from the animal, fusing the B cells with a myeloma cells to produce hybridomas, obtaining the monoclonal antibodies from the hybridomas.

Antibodies to the outer membrane proteins of *E. chafeensis* and *E. canis* are useful research tools for identifying cells, particularly monocytes, infected with *E. chafeensis* or *E. canis* and for purifying the corresponding outer membrane protein of *E. chafeensis* or *E. canis* from partially purified preparations by affinity chromatography. Such antibodies are also useful for identifying bacterial colonies, particularly colonies of genetically-engineered bacteria, that are expressing the major outer membrane protein.

Diagnostic Method

The present invention also provides a method for detecting antibodies to the *E. chafeensis* or *E. canis* in a sample of a bodily fluid from a patient. The method comprises providing an isolated outer membrane protein of *E. chafeensis* or *E. canis*, particularly a recombinant form of the isolated protein, contacting the outer membrane protein or polypeptide with a sample taken from the patient; and assaying for the formation of a complex between the outer membrane protein or polypeptide and antibodies in the sample. For ease of detection, it is preferred that the isolated protein or polypeptide be attached to a substrate such as a column, plastic dish, matrix, or membrane, preferably nitrocellulose. The sample may be a tissue or a biological fluid, including urine, whole blood, or exudate, preferably serum. The sample may be untreated, subjected to precipitation, fractionation, separation, or purification before combining with the isolated protein or peptide. Interactions between antibodies in the sample and the isolated protein or peptide are detected by radiometric, colorimetric, or fluorometric means, size-separation, or precipitation. Preferably, detection of the antibody-outer membrane protein complex is by addition of a secondary antibody that is coupled to a detectable tag, such as for example, an enzyme, fluorophore, or chromophore. Formation of the complex is indicative of the presence of anti-*E. chafeensis* or anti-*E. canis* antibodies,

either IgM or IgG, in the patient. Thus, the method is used to determine whether a patient is infected with *E. chaffeensis* or *E. canis*.

Preferably, the method employs an enzyme-linked immunosorbent assay (ELISA) or a Western immunoblot procedure. Such methods are relatively simple to perform and do not require special equipment as long as membrane strips are coated with a high quality antigen. Accordingly, it is more advantageous to use a recombinant form of the outer membrane protein of *E. chaffeensis* or *E. canis* since such proteins, typically, are more pure and consistent in quality than a purified form of such protein.

Immunogenic Composition

The present invention also relates to immunogenic compositions comprising one or more of the isolated outer membrane proteins of *E. chaffeensis* and a pharmaceutically acceptable adjuvant and to immunogenic compositions comprising an isolated P30 protein of *E. canis* and a pharmaceutically acceptable adjuvant, which, preferably, enhances the immunogenic activity of the outer membrane protein in the host animal.

Preparation of a Polynucleotide which Encodes OMP-1(P28)

A. Isolation of the Outer Membrane Proteins

E. chaffeensis Arkansas strain and *E. canis* Oklahoma strain were cultivated in the DH82 dog macrophage cell line and purified by Percoll density gradient centrifugation. Purified ehrlichiae (100 µg) were suspended with 10 mM sodium phosphate buffer, pH 7.4, containing 0.1% Sodium N-lauroyl sarcosine (Sarkosyl) [Sigma, St. Louis, MO], 50 µg/ml each DNase I (Sigma) and RNase A (Sigma), and 2.5 mM MgCl₂. After incubation at 37° for 30 min, the sample was separated by centrifugation at 10,000 x g for 1 h into the soluble supernatant and the insoluble precipitate. The insoluble pellet was resuspended 2 to 3 times with 0.1% Sarkosyl and centrifuged. The final pellet was analyzed by sodium dodecyl sulfate (SDS)-polyacrylamide gel electrophoresis (PAGE) and by electron microscopy.

Transmission electron microscopy revealed that the purified ehrlichial fraction consists of a mixture of electron dense and light forms of *E. chaffeensis* with slight disintegration of inner membrane. Ehrlichiae were not surrounded with the host inclusion membrane. Various sizes of membrane vesicles (< 1 µm) without significant ribosomes or nuclear materials were observed in the Sarkosyl-insoluble fraction from the organism. Succinic dehydrogenase (inner membrane marker enzyme of gram negative bacteria) activities were at less than the detection limit (1 n moles / min / mg of protein) in the Sarkosyl-insoluble fraction compared to approximately 10 n moles / min / mg of protein in the Percoll-purified organisms, suggesting that the insoluble fraction primarily consisted of the outer membrane of *E. chaffeensis*.

Analysis of the Sarkosyl-soluble, and insoluble fraction of *E. chaffeensis* by SDS-PAGE suggested that proteins of 30-kDa range in the insoluble fraction represent the major outer membrane proteins of this organism. Analysis of the Sarkosyl-soluble, and insoluble fraction of *E. canis* by SDS-PAGE suggested that proteins of 30-kDa range in the insoluble fraction represent the major outer membrane proteins of this organism also. *E. canis* was

antigenically cross reactive with *E. chafeensis*. These findings indicate that the 30-kDa range proteins represent the major outer membrane proteins of these two *Ehrlichia* spp.

To improve resolution of the outer membrane proteins, proteins in the Sarkosyl-insoluble pellet prepared from 400 µg of purified *E. chafeensis* were separated by a reversed-disc continuous (Rd) SDS-PAGE (2.5-cm-long 17% gel on top of 11-cm-long 12% gel). At least five proteins of 30-kDa range in *E. chafeensis* (P23, P25, P27, P28, and P29) were resolved from the Sarkosyl-insoluble proteins.

B. Cloning and sequencing of the *p28* gene

The portion of the membrane containing bound proteins was excised and analyzed with an Applied Biosystems protein sequencer (Model 470). The N-terminal amino acid sequence of P28 was determined as D P A G S G I N G N F Y S G K Y M P, SEQ IN NO _____. Based on 6th to 12th amino acids of this sequence, a forward primer, FECH1, having the sequence: 5'-
CGGGATCCGAATTCGG(A/T/G/C)AT(A/T/C)AA(T/C)GG(A/T/G/C)AA(T/C)TT(T/C)TA-3'. SEQ ID NO _____. was designed. Amino acids at the 1 to 5 positions of the N terminus of P28 were not included in this primer design. For insertion into an expression vector, a 14-bp sequence (underlined) was added at the 5' end of primer to create an *Eco*RI and a *Bam*HI site. The reverse primer, RECH2, which includes a *Nol*I site at the 5' end for ligation into an expression vector had the sequence: 5'-AGCGGCCGCTTA(A/G)AA(T/C)A(C/G) (A/G)AA (C/T)CT T(C/G)C TCC-3'. SEQ ID NO _____.

Genomic DNA of *E. chafeensis* was isolated from purified organisms. PCR amplification with FECH1 and RECH2 primers was performed using a Perkin-Elmer Cetus DNA Thermal Cycler (model 480). A 0.8-kb amplified product was cloned in the pCRII vector of a TA cloning kit, as described by the manufacturer (Invitrogen Co., San Diego, CA). The clone obtained was designated pCRII_p28. Both strands of the inserted DNA were sequenced by a dideoxy-termination method with an Applied Biosystems 373A DNA sequencer.

The 0.8-kb DNA fragment, cloned in pCRII_p28, had an open reading frame (ORF) of 756 bp encoding a 251-amino acid recombinant protein (including both PCR primer regions) with a molecular mass of 27,685 Da. The nucleotide sequence of the open reading frame, SEQ ID NO: ___, and the amino acid sequence of the polypeptide of the OMP-1 protein, SEQ ID NO ___, are shown in Figs ____ and _____, respectively.

A DNA fragment comprising the *p30* gene was prepared in a similar manner, i.e., by PCR amplification of genomic DNA of *E. canis* with the FECH1 and RECH2 primers.

Preparation of Polynucleotides which encode OMP 1A, OMP1B, OMP1-C, OMP-1D, OMP-1F, and OMP1-E

A. Southern blot analysis. Genomic DNA extracted from the purified *E. chafeensis* (200 ng each) was digested with restriction endonucleases, electrophoresed, and transferred to Hybond-N⁺ nylon membrane (Amersham, Arlington Heights, IL), by a standard method. The 0.8-kb *p28* gene fragment from the clone pCRII_p28 was labeled with [α -³²P]dATP by the random primer method using a kit (Boehringer Mannheim, Indianapolis, IN) and the labeled fragment was used as a DNA probe. Hybridization was performed at 60°C in rapid hybridization buffer (Amersham) for 20 h. The nylon sheet was washed in 0.1 x SSC (1 x SSC containing 0.15M sodium chloride and

0.015M sodium citrate) with 1% SDS at 55°C and the hybridized probes were exposed to Hyperfilm (Amersham) at -80°C.

Genomic Southern blot analysis with several restriction enzymes resulted in one or more DNA fragment(s) of *E. chaffeensis* which hybridized to ³²P-labeled *p28* gene probe. The restriction enzymes used did not cut within the *p28* gene portion of the pCRII*p28* insert. *Xba* I, *Bgl* II, and *Kpn* I produced two bands, *Spe* I generated three bands, and *Eco*R V and *Pst* I produced multiple bands with different densities. *Eco*R I generated a broad band of 2.5 to 4kb. These *p28* homologous genes are designated as *omp-1* (outer membrane protein-1) family.

B. Cloning and sequencing of genomic copies of *E. chaffeensis* *p28* gene. The *Eco*R I and *Pst* I fragments of DNA, detected by genomic Southern blot analysis as described above, were inserted into pBluescript II KS (+) vectors, and the recombinant plasmids were introduced into *E. coli* DH5α. Using the colony hybridization method with the ³²P-labeled *p28* gene probe, four positive clones were isolated from the transformant. The positive clones were designated pEC2.6, pEC3.6, pPS2.6, and pPS3.6. These contained the ehrlichial DNA fragments of 2.6-kb (*Eco*R I), 3.6 kb (*Eco*R I), 2.6 kb (*Pst* I), and 3.6 kb (*Pst* I), respectively. The inserts of the clones pEC3.6 and pPS2.6 overlapped as shown in Fig. _____. The overlapping area was further confirmed by PCR of *E. chaffeensis* genomic DNA with two pairs of primer sets interposing the junctions of the four clones. The 1.1- to 1.6-kb DNA fragments of *Hind*III-*Hind*III, *Hind*III-*Eco*RI, or *Xba*I-*Eco*RI in the pEC2.6 and pEC3.6 were subcloned for sequencing. DNA sequencing was performed with suitable synthetic primers by dideoxy-termination method as described above.

Four DNA fragments from 2.6 to 3.6 kb were cloned from the *Eco*RI-digested and the *Pst*I-digested genomic DNA of *E. chaffeensis* by colony hybridization with radiolabeled *p28* gene probe. The inserted DNA of the two recombinant clones, pEC3.6 and pPS2.6, were overlapped as shown in Fig. 7. Sequencing revealed one 5'-truncated ORF of 243 bp (designated *omp-1A*) and five complete ORF of 836-861 bp (designated *omp-1B* to *omp-1F*), which are tandemly-arrayed and are homologous to the *p28* gene (but are not identical), in the ehrlichial genomic DNA of 6,292 bp. The intergenic spaces were 581 bp between *omp-1A* and *omp-1B* and 260-308 bp among others. Putative promoter regions and ribosome-binding sites were identified in the noncoding regions.

Sequence analysis and GenBank accession number.

Nucleotide sequences were analyzed with the DNASIS program (Hitachi Software Engineering Co., Ltd., Yokohama, Japan). A homology search was carried out with databases of the GenBank, Swiss Plot, PDB and PIR by using the software basic local alignment search tool in the BLAST network service (the National Center for Biotechnology Information, Bethesda, MD). Phylogenetic analysis was performed by using the PHYLIP software package (version 3.5). An evolutional distance matrix, generated by using the Kimura formula in the PROTDIST, was used for construction of a phylogenetic tree by using the unweighted pair-group method analysis (UPGMA) (Felsenstein, J. 1989. PHYLIP-phylogeny inference package (version 3.3). Cladistics 5:164-166). The data were also examined using parsimony analysis (PROTPARS in PHYLIP). A bootstrap analysis was carried out to investigate the stability of randomly generated trees by using SEQBOOT and CONSENSE in the same package. The nucleotide sequence of the *p28* gene and its gene copies has been assigned GenBank accession numbers U72291 and AF021338, respectively.

Proteins of the *E. chaffeensis* *omp-1* Family.

Five complete *omp-1* gene copies (*omp-1B* to *omp-1F*) encode 279 to 287-amino acid proteins with molecular masses of 30,320 - 31,508 Da. *Omp-1A* encodes an 82-amino acid partial protein (9,243 Da) which lacks the N-terminal region. The 25-amino acid sequence at the N-terminus of OMP-1B to OMP-1F (encoded in *omp-1B* to *omp-1F*) is predicted to be a signal peptide because three carboxyl-terminal amino acids of the signal peptides (Ser-X-Ala in OMP-1B, Leu-X-Ser for OMP-1C, and Ser-X-Ser for OMP-1D and OMP-1F) are included in the preferred amino acid sequence of signal peptidase at the processing sites proposed by Oliver. The calculated molecular masses of the mature OMP-1B to OMP-1F from the predicted amino acid sequences are 28,181 Da for OMP-1B, 27,581 Da for OMP-1C, 28,747 Da for OMP-1D, 27,776 Da for OMP-1E, and 27,933 Da for OMP-1F. The estimated isoelectric points are 4.76-5.76 in the mature OMP-1B to OMP-1F. An amino acid sequence in *omp-1F* gene (the 80th to 94th amino acids) was identical to the N-terminal amino acid sequences of *E. chaffeensis* native P23 protein as determined chemically, which indicates that P23 is derived from the *omp-1F* gene. Amino acid sequences identical to the N-terminal sequences of P25, P27, and P29 were not found in those from *omp-1* gene copies cloned in this study.

Alignment of predicted amino acid sequences of the *E. chaffeensis* OMP-1 family and *Cowdria ruminantium*, revealed substitutions or deletions of one or several contiguous amino acid residues throughout the molecules. The significant differences in sequences among the aligned proteins are seen in the regions indicated SV (semivariable region) and HV (hypervariable region) 1 to 3 in Fig 3I. Computer analysis for hydropathy revealed that protein molecules predicted from all *omp-1* gene copies contain alternative hydrophilic and hydrophobic motifs which are characteristic of transmembrane proteins. The HV1 and HV2 were found to locate in the hydrophilic regions.

The amino acid sequences of 5 mature proteins without signal peptides (OMP-1C to OMP-1F and a P28) were similar to one another (71-83%) but the sequence of OMP-1B was dissimilar to those of the 5 proteins (45-48%). The amino acid sequences of the 5 proteins showed an intermediate degree of similarity with that of *C. ruminantium* MAP-1 (59-63%), but the similarity between that of the OMP-1B and the *C. ruminantium* MAP-1 was low (45%). These relations are shown in a phylogenetic tree which was obtained based on the amino acid sequence alignment by UPGMA method in the PHYLIP software package (Fig. 10). Three proteins (P28, OMP-1D, and OMP-1F) and two proteins (OMP-1C and OMP-1E) formed two separate clusters. The OMP-1B was located distantly from these two clusters. The *C. ruminantium* MAP-1 was positioned between the OMP-1B and other members in the OMP-1 family.

Preparation of a Recombinant form of OMP-1 and P30

The 0.8-kb *p28* gene was excised from the clone pCRII_p28 by *Eco*RI-*Not*I double-digestion, ligated into *Eco*RI-*Not*I sites of a pET 29a expression vector, and amplified in *Escherichia coli* BL21 (DE3)pLysS (Novagen, Inc., Madison, WI). The clone (designated pET29_p28) produced a fusion protein with a 35-amino acid sequence

carried from the vector at the N terminus. The amino acid sequence of the OMP-1 portion of the fusion protein is depicted in Fig. 1.

An expression vector comprising the p30 gene was used to prepare the recombinant form of P30.

The following examples are for purposes of illustration only and are not intended to limit the scope of the claims which are appended hereto.

Preparation of anti rP28 (anti-OMP1) antibody

The (r) P28 antigen was prepared by excising the gel band corresponding to the rP28 in SDS-PAGE, mincing the band in phosphate-buffered saline (PBS), pH 7.4, and mixing with an equal volume of Freund's incomplete adjuvant (Sigma). The rP28 mixture (1 mg of protein each time) was subcutaneously injected into a rabbit every 2 weeks four times. A serum sample was collected from the rabbit to provide the anti-rP28 antibody.

The anti-rP28 antibody was examined by western immunoblots analysis. The results indicated that the rabbit anti-rP28 antibody recognized not only rP28 (31 kDa) and P28, but also P29 and P25 of *E. chaffeensis* and P30 of *E. canis*. These results indicate that P28 shares antigenic epitopes with P25 and P29 in *E. chaffeensis* and P30 of *E. canis*.

Example 1. Assaying for the presence of anti-OMP-1 antibody in a Patient

Convalescent-phase serum from a patient with clinical signs of human ehrlichiosis was used. Western blot analyses using the rP28 protein as antigen was performed with 1:1,000 dilutions of this serum. Alkaline phosphatase-conjugated affinity-purified anti-human, anti-rabbit or anti-mouse immunoglobulin G (Kirkegaard & Perry Laboratories, Inc., Gaithersburg, MD) were used at a 1:1,000 or 1:2,000 dilution as secondary antibodies. Results indicated that serum from a patient with clinical signs of human ehrlichiosis reacted strongly to rP28 (31 kDa).

Example 2. Assaying for the presence of anti-OMP-1 antibody in a Patient

Convalescent-phase serum from a patient with clinical signs of human ehrlichiosis was reacted with the rP30 protein of *E. canis* as described in Example 1. The serum reacted strongly to rP30. These results indicate the rP30 is useful for diagnosing an infection with *E. chaffeensis* in human patients.

Example 3. Identifying *E. chaffeensis*-infected cells using anti-rP 28 antibody

E. chaffeensis-infected DH82 cells were sonicated and centrifuged at 400 x g for 10 min. The supernatant was then centrifuged at 10,000 x g for 10 min to obtain ehrlichia-enriched pellet. The pellet was resuspended and incubated with rabbit anti-rP28 antibody or normal rabbit serum (1:100 dilution) at 37°C for 1h in PBS containing 1% bovine serum albumin (BSA-PBS). After washing, the ehrlichiae was incubated with gold-conjugated protein G (20 nm, Sigma) at 1:30 dilution for 1 h at room temperature in BSA-PBS. After washing again, the specimen was fixed with 1.25% formaldehyde, 2.5% glutaraldehyde, and 0.03% trinitrophenol in 0.1 M cacodylate buffer (pH 7.4) for 24h and postfixed in 1% osmium-1.5% potassium ferricyanide for 1 h (34). The section was then embedded in

PolyBed 812 (Polysciences, Warraington, Pa). The specimen was ultrathin sectioned at 60 nm, stained with uranyl acetate and lead citrate, and observed with a Philips 300 transmission electron microscope at 60 kV.

Transmission immunolectron microscopy with colloidal gold-conjugated protein G and rabbit anti-rP28 antibody revealed gold particles bound to *E. chaffeensis* surface. The distribution of the particles was random, close to the surface, and appeared as if almost embedded in the membrane, suggesting that the antigenic epitope protrudes very little from the lipid bilayer. Nonetheless, the antigenic epitope was surface-exposed, and thus, could be recognized by rabbit anti-rP28 antibody. No gold particles were observed on host cytoplasmic membrane or *E. chaffeensis* incubated with normal rabbit serum.

Example 4. Immunization of mice and *E. chaffeensis* challenge.

The rP28 band in SDS-PAGE was excised, minced, and mixed with an equal volume of Freund's incomplete or complete adjuvant. Nine BALB/c male mice (6 weeks old) were divided into two groups. Five mice were intraperitoneally immunized a total of four times at 10-day intervals; twice with a mixture of the minced gel with the rP28 (30 to 40 µg of protein per mouse each time) and incomplete adjuvant, and twice with a mixture of the recombinant protein (the same amount as before) and complete adjuvant. Four mice were intraperitoneally injected with a mixture of the minced gel without protein and the respective adjuvants. For ehrlichia-challenge, approximately 1×10^7 DH82 cells heavily-infected with *E. chaffeensis* were disrupted by sonication in serum-free DMEM (GIBCO-BRL) and centrifuged at 200 x g for 5 min. The supernatant was diluted to a final volume of 5 ml, and 0.3 ml was inoculated intraperitoneally into each mouse 10 days after the last immunization. Before challenge, all 5-immunized mice had a titer of 1:160 against *E. chaffeensis* antigen by IFA and all 4-nonimmunized mice were negative.

At day 5 post-challenge, approximately 1 ml of blood was collected in an EDTA tube from each mouse and protection was assessed by PCR detection of *E. chaffeensis* 16S rDNA in the buffy coat of the collected blood. *E. chaffeensis* could not be reisolated in cell culture at day 10 postinfection. Day 5 post challenge is the optimum time at which establishment of ehrlichial infection can be examined by PCR without the influence of residual DNA from the ehrlichiae used as the challenge before the spontaneous clearance of organisms take place. The *E. chaffeensis*-specific DNA fragment was observed in all nonimmunized mice but not in any immunized mice, indicating that immunization of rP28 apparently protects mice from ehrlichial infection and indicating that the P28 is a potential protective antigen.

Example 5 Assaying for the presence of anti-P30 antibody in Dogs

The rP30 protein was used as an antigen in a Western immunoblot analysis and dot blot analysis to detect the presence of antibody to *E. canis* in serum from *E-canis* infected dogs. The results of the Western immunoblot analysis indicated that reactivity of the sera with rP30 was stronger than the reactivity that was observed when purified *E.canis* was used as antigen. The results of the dot blot assay indicated that rP30 is a useful and sensitive tool for serodiagnosis of canine ehrlichiosis.

CLAIMS

What is claimed is:

1. An isolated polynucleotide encoding an outer membrane protein of *E. chaffeensis* or an immunogenic fragment thereof, wherein the outer membrane protein is selected from the group consisting of OMP-1, OMP-1B, OMP-1C, OMP-1D, OMP-1E, OMP-1F, OMP-1R, OMP-1S, OMP-1T, OMP-1U OMP-1V, OMP-1W OMP-1X OMP-1Y, and OMP-1Z.
2. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an OMP-1 protein comprising a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 3B, SEQ. ID NO ____.
3. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an OMP-1B protein comprising a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 4B, SEQ. ID NO ____.
4. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an OMP-1C protein comprising a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 5B, SEQ. ID NO ____.
5. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an OMP-1D protein comprising a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 6B, SEQ. ID NO ____.
6. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an OMP-1E protein comprising a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 7B, SEQ. ID NO ____.
7. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an OMP-1F protein comprising a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 8B, SEQ. ID NO ____.
8. The isolated polynucleotide of claim 1 wherein the polynucleotide encodes an immunogenic fragment of the OMP-1 protein, said fragment comprising a sequence which is at least 85% homologous to the amino acid sequence extending from amino acid 6 through amino acid 251 as shown in Fig. 1, SEQ. ID NO ____.
9. An isolated polynucleotide encoding an outer membrane protein of *E. canis* or an immunogenic fragment thereof, wherein the outer membrane protein is selected from the group consisting of P30, P30-A, P30-1, P30-2, P30-3, P30-4, P30-5, P30-6, P30-7, P30-8, P30-9, P30-10.
10. The isolated polynucleotide of claim 9 wherein said P30 protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 19 SEQ ID NO ____.
11. An isolated outer membrane protein of *E. chaffeensis* or an immunogenic fragment thereof, wherein said protein is selected from the group consisting of OMP-1, OMP-1B, OMP-1C, OMP-1D, OMP-1E, OMP-1F, OMP-1R, OMP-1S, OMP-1T, OMP-1U OMP-1V, OMP-1W OMP-1X OMP-1Y, and OMP-1Z.
12. The isolated OMP-1 protein of claim 11, wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 4B, SEQ. ID NO ____.
13. The isolated OMP-1B protein of claim 11 wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 5B, SEQ. ID NO ____.
14. The isolated OMP-1C protein of claim 11 wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 6B, SEQ. ID NO ____.

15. The isolated OMP-1D protein of claim 11 wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 7B, SEQ. ID NO ____.
16. The isolated OMP-1E protein of claim 11 wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 8B, SEQ. ID NO ____.
17. The isolated OMP-1F protein of claim 11 wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig. 9B, SEQ. ID NO ____.
18. The isolated immunogenic fragment of the OMP-1 protein of claim 11, said fragment comprising a sequence which is at least 85% homologous to the amino acid sequence extending from amino acid 6 through amino acid 251 as shown in Fig. 1, SEQ. ID NO ____.
19. An isolated outer membrane protein of *E. canis* or an immunogenic fragment thereof, wherein the outer membrane protein is selected from the group consisting of P30, P30-A, P30-1, P30-2, P30-3, P30-4, P30-5, P30-6, P30-7, P30-8, P30-9, P30-10.
20. The isolated P-30 protein of claim 19 wherein said protein comprises a sequence which is at least 85% homologous to the amino acid sequence shown in Fig 19, SEQ ID NO ____.
21. A method for diagnosing an infection with *E. chaffeensis* in a patient comprising the steps of:
 - (a) providing a serum sample from the patient;
 - (b) providing an outer membrane protein selected from the group consisting of a protein of claim 11, a protein of claim 19, and mixtures thereof;
 - (c) contacting the serum sample with the outer membrane protein; and
 - (d) assaying for the formation of a complex between antibodies in the serum sample and the outer membrane protein, wherein formation of said complex is indicative of infection with *E. chaffeensis*.
22. A method for diagnosing an infection with *E. canis* in a Canidae patient comprising the steps of:
 - (a) providing a serum sample from the patient;
 - (b) providing an outer membrane protein of claim 19;
 - (c) contacting the serum sample with the outer membrane protein; and
 - (d) assaying for the formation of a complex between antibodies in the serum sample and the outer membrane protein, wherein formation of said complex is indicative of infection with *E. canis*.

220p1 pridr →
 GGATATAATGGGAATTTCATCATCGAAATACATCCAACTGCTTGCATTTGG
 9 P A D O O G I N G N F Y I D O G E Y H P D A D V R G
 230
 GATATCTCTGTTAAAGGAGAAAGAAATACACAGTGTGAGTTTGTGAACTGACCAAAATTTGGGACGGAGGCGCATATTCACACTCC
 V P D A E B D H T T V G V P D L E K O N H B G D A I O N D B
 150
 95
 CCAAACGATGATTCATCTCAAAATATTCAATTAAATATGAAACCAACCCGTTTTAGGTTTGCAGGAGCTATGGTACTCAATG
 P H D V F T V G H Y S P R E T E H P F L G F A G A I G Y S M
 200
 205
 GATGTTCCAAAGATAGAGCTGTGAACTTCATGAAACATTTTATGAAAAATCAAGGTAAACAAATGAAATGAGGCAATAGATAT
 D G F R I E E B V G Y E T F D V R H Q G H N H T K H A R R Y
 130
 115
 TUTGCTCTATCCATAACTCAGCGCAGACATGAGTACTGCAAGTAAATTTTGTCTTCTAAAGATGAGGATTACTGCAATATCA
 C A L B R H S A A D H S G A B D N F V P L R D B G L L D I S
 420
 145
 TTATGCTGAAACGATGCTATGCGTGTAGGCGAAAGGCAACCTTTCTCCCTTAAATGCGCAAGTATCGGACTGTATTTATCC
 V H L D A C T D V V G E G I P P S P Y I C A G I G T D L V S
 510
 175
 ATGGTTGAACTACAAATCCATAAAATTCCTTACCGAGGAAAGTTAGGTTTACGCTACTCTATANGCCAGAGCTCTGTGTTTATGGT
 H P E A T H P K I O Y Q G R L G L S T S K S P R A D V F I G
 600
 205
 GGGCACTTCATAAGGTTAATGGGAGACGAAATTAGAGATATTCCTACTATAATACCTACTGCACTGCAACTGGCAGGAAAGAAATAC
 G H P H R V I G H E P R D I P T I I P T G T O S T L A G E G O N Y
 690
 235
 CCTGAACTAAATACGGATGATGCGACTTGTGAACTGAGGAGGAGGTTATTCCTAA
 P A I V I L D V C H F G I E L G G R F V P * ←
 756
 256
 r20p1 pridr

Fig. 1




Fig. 2

2/31

10	20	30	40	50	60
ATGAATTACA	AAAAAGTTTT	CATAACAAGT	GCATTGATAT	CATTAATATC	TTCTCTACCT
70	80	90	100	110	120
GGAGTATCAT	TTTCCGACCC	AGCAGGTAGT	GGTATTAACG	GTAATTCTCA	CATCAGTGGA
130	140	150	160	170	180
AAATACATGC	CAAGTGCTTC	GCATTTGGA	GTATTCTCTG	CTAAGGAAGA	AAGAAATACA
190	200	210	220	230	240
ACAGTTGGAG	TGTTTGGACT	GAAGCAAAAT	TGGGACGGAA	GCGCAATATC	CAACTCCTCC
250	260	270	280	290	300
CCAAACGATG	TATTCACTGT	CTCAAAATTAT	TCATTTAAT	ATGAAAACAA	CCCGTTTTTA
310	320	330	340	350	360
GGTTTGCAAG	GAGCTATTGG	TTACTCAATG	GATGGTCCAA	GAATAGAGCT	TGAAGTATCT
370	380	390	400	410	420
TATGAAACAT	TTGATGTAAA	AAATCAAGGT	ACAAATTATA	AGAATGAAGC	ACATAGATAT
430	440	450	460	470	480
TGTGCTCTAT	CCCATAACTC	AGCAGCAGAC	ATGAGTAGTG	CAAGTAATAA	TTTTGTCTTT
490	500	510	520	530	540
CTAAAAAAATG	AAGGATTACT	TGACATATCA	TTTATGCTGA	ACGCATGCTA	TGACGTAGTA
550	560	570	580	590	600
GGCGAAGGCA	TACCTTTTTC	TCCTTATATA	TGCGCAGGTA	TCGGTACTGA	TTTAGTATCC
610	620	630	640	650	660
ATGTTTGAAG	CTACRAATCC	TAAAATTCT	TACCAAGGAA	AGTTAGGTTT	AAGCTACTCT
670	680	690	700	710	720
ATAAGCCCAG	AAGCTCTGT	GTATTATTGGT	GGGCACITTC	ATAAGGTAAT	AGGGAAACGAA
730	740	750	760	770	780
TTTAGAGATA	TTCCTACTAT	AATACCTACT	GGATCAACAC	TTGCAGGAAA	AGGAAACTAC
790	800	810	820	830	840
CCTGCAATAG	TAATACTGGA	TGTATGCCAC	TTTGGAAATAG	AACTTGGAGG	AAGGTTTGTAA
850	860	870	880	890	900
TTCTAA.....

Fig. 3A

10	20	30	40	50	60
MNYKKVFIITS	ALISLISSLP	GVSFSDPAGS	GINGNFYISG	KYMP SASHFG	VFSAKEERNT
70	80	90	100	110	120
TVGVFGLKQN	WDGSAISNSS	PNDVFTVSNY	SFKYENNPF	GFAGAIGYSM	DGPRIELEVS
130	140	150	160	170	180
YETFDVKNQG	NNYKNEAHRY	CALSHNSAAD	MSSASNNFVF	LKN EGLLDIS	FMLNACYDVV
190	200	210	220	230	240
GEGIPFSPYI	CAGIGTDLVS	MFEATNPKIS	YQKLGLSYS	ISPEASVFIG	GHFKVIGNE
250	260	270	280	290	300
FRDIPTIIPT	GSTLAGKGNY	PAIVILDVCH	FGIEJ-GGRFV	F.....

Fig. 3B

10	20	30	40	50	60
ATGAATTACA	AGAAAATTT	TGTAAGCAGT	GCATTAATTT	CATTAATGTC	AATCTTACCT
70	80	90	100	110	120
TACCAATCTT	TTGCAGATCC	TGTAACCTCA	AATGATAACAG	GAATCAACGA	CAGCAGAGAA
130	140	150	160	170	180
GGCTTCTACA	TTAGTGTAAA	GTATAATCCA	AGCATATCAC	ACTTCAGAAA	ATTCTCAGCT
190	200	210	220	230	240
GAAGAAGCTC	CCATCAATGG	AAATACTTCT	ATCACTAAAAA	AGGTTTCGG	GCTGAAAAAA
250	260	270	280	290	300
GACGGGAGATA	TAGCACAATC	TGCGAATTT	AACAGGACAG	ATCCAGCCCT	CGAGTTTCAG
310	320	330	340	350	360
AATAACCTAA	TATCAGGATT	CTCAGGAAGT	ATTGGTTATG	CTATGGATGG	GCCAAGAATA
370	380	390	400	410	420
GAACATTGAAG	CTGCATACCA	AAAATTGAT	GCAAAAAATC	CTGACAACAA	TGACACTAAT
430	440	450	460	470	480
AGCGGTGACT	ACTATAAATA	CTTGGACTA	TCTCGTGAAG	ACGCAATAGC	AGATAAGAAA
490	500	510	520	530	540
TATGTTGTCC	TTAAAAATGA	AGGCATCACT	TTTATGTCAT	TAATGGTTAA	CACTTGCTAT
550	560	570	580	590	600
GACATTACAG	CTGAAGGAGT	ACCTTTCATA	CCGTATGCAT	GTGCAGGTGT	AGGAGCAGAC
610	620	630	640	650	660
CTTATAAACG	TATTAAGGA	TTTTAATTAA	AAATTCTCAT	ACCAAGGGAA	AATAGGTATT
670	680	690	700	710	720
AGCTATCCAA	TCACACCAAGA	AGTTTCCGCT	TTTATTGGAG	GATACTACCA	CGGAGTTATA
730	740	750	760	770	780
GGAAAATAATT	TTAACAAAAT	ACCTGTAATA	ACACCTGTAG	TATTAGAAGG	AGCTCCTCAA
790	800	810	820	830	840
ACCACATCTG	CGCTAGTAAC	TATTGACACT	GGATACTTG	GCGGAGAAGT	TGGAGTAAGG
850	860	870	880	890	900
TTCACCTTCT AG.....					

Fig. 4A

10	20	30	40	50	60
MNYKKIFVSS	ALISILMSILP	YQSFADPVTS	NDTGINDSRE	GFYISVKYNP	SISHFRKFSA
70	80	90	100	110	120
EEAPINGNTS	ITKKVFGLKK	DGDIAQSANF	NRTDPALEFQ	NNLISGFSGS	IGYAMDGPRI
130	140	150	160	170	180
ELEAAAYQKFD	AKNPDNNDTN	SGDYYKYFGL	SREDAIADKK	YVVLKNEGIT	FMSILMVNTCY
190	200	210	220	230	240
DITAEGVPF	PYACAGVGAD	LINVFKDFNL	KPSYQGKIGI	SYPITPEVSA	FIGGYYHGVI
250	260	270	280	290	300
GNNFNKIPVI	TPVVLEGAPQ	TTSALVTIDT	GYFGGEVGVR	FTF.....

Fig. 4B

10	20	30	40	50	60
ATGAACTGCA	AAAAATTTT	TATAACAAC	GCATTGGCAT	TGCCAATGTC	TTTCTTACCT
70	80	90	100	110	120
GGAATATTAC	TTTCTGAACC	AGTACAAGAT	GACAGTGTGA	GTGGCAATT	CTATATTAGT
130	140	150	160	170	180
GGCAAGTACA	TGCCAAGTGC	TTCTCATTT	GGAGTTTCT	CTGCCAAAGA	AGAAAAAAAT
190	200	210	220	230	240
CCTACTGTCG	CGTTGTATGG	TTTGAAACAA	GATTGGAACG	GTGTTAGTGC	TTCAAGTCAT
250	260	270	280	290	300
GCTGATGCGG	ACTTTAATAA	CAAAGGTTAT	TCTTTTAAAT	ACGAAAACAA	TCCATTTCTA
310	320	330	340	350	360
GGTTTGCAG	GAGCTATTGG	TTATTCAATG	GGTGGTCCAA	GAATAGAGTT	TGAAGTGTCC
370	380	390	400	410	420
TATGAAACAT	TTGACGTGAA	AAATCAAGGT	GGTAATTACA	AAAATGATGC	TCACAGATAC
430	440	450	460	470	480
TGTGCCTTAG	ATCGTAAAGC	AAGCAGCACT	AATGCCACAG	CTAGTCACTA	CGTGCTACTA
490	500	510	520	530	540
AAAAATGAAG	GACTACTTGA	TATATCACTT	ATGTTGAATG	CATGCTATGA	CGTAGTAAGT
550	560	570	580	590	600
GAAGGAATAC	CTTTCTCTCC	TTACATATGT	GCAGGTGTTG	GTACCGATT	AATATCCATG
610	620	630	640	650	660
TTTGAAGCTA	AAACCCCTAA	AATTCTTAT	CAAGGAAAGT	TAGGTTTGAG	TTACTCTATA
670	680	690	700	710	720
AACCCAGAAG	CTTCTGTCTT	TGTTGGTGG	CATTTTCATA	AAGTTGCAGG	TAATGAATTC
730	740	750	760	770	780
AGGGACATTT	CTACTCTTAA	AGCGTTTGCT	ACACCATCAT	CTGCAGCTAC	TCCAGACTTA
790	800	810	820	830	840
GCAACAGTAA	CACTGAGTGT	GTGTCACTTT	GGAGTAGAAC	TTGGAGGAAG	ATTTAACTTC
850	860	870	880	890	900
TAA.....					

Fig. 5A

10	20	30	40	50	60
MNCKKFFITT	ALALPMSFLP	GILLSEPVQD	DSVSGNFYIS	GKYMP SASHF	GVFSAKEEKN
70	80	90	100	110	120
PTVALYGLKQ	DWNGVSASSH	ADADFNNKGY	SFKYENNPF	GFAGAIGYSM	GGPRIEFEV
130	140	150	160	170	180
YETFDVKNQG	GNYKNDAHRY	CALDRKASST	NATASHYVLL	KNEGLLDISL	MLNACYDVVS
190	200	210	220	230	240
EGIPFSPYIC	AGVGTDLISM	FEAINPKISY	QGKLGLSYSI	NPEASVFVGG	HFHKVAGNEF
250	260	270	280	290	300
RDISTLKAF	TPSSAATPDL	ATVTL SVCHF	GVELGGRNF

Fig. 5B

10	20	30	40	50	60
ATGAACTGCG	AAAAATTTT	TATAACAAC	GCATTAACAT	TACTAATGTC	CTTCTTACCT
70	80	90	100	110	120
GGAATATCAC	TTTCTGATCC	AGTACAGGAT	GACAACATTA	GTGGTAATT	CTACATCACT
130	140	150	160	170	180
GGAAAGTATA	TGCCAAGCGC	TTCGCATT	GGAGTTTTT	CTGCCAAGGA	AGAAAGAAAT
190	200	210	220	230	240
ACAACAGTTG	GAGTATTG	AATAGAGCAA	GATTGGGATA	GATGTGTAAT	ATCTAGAAC
250	260	270	280	290	300
ACTTTAACCG	ATATATTAC	CGTTCCAAAT	TATTCATT	AGTATGAAA	TAATCTATT
310	320	330	340	350	360
TCAGGATTG	CAGGAGCTAT	TGGCTACTCA	ATGGATGGCC	CAAGAATAGA	GCTTGAAGTA
370	380	390	400	410	420
TCTTATGAAG	CATTCGATGT	AAAAATCAA	GGTAACAATT	ATAAGAACGA	AGCACATAGA
430	440	450	460	470	480
TATTATGCTC	TGTCCCACATCT	TCTCGGCACA	GAGACACAGA	TAGATGGTGC	AGGCAGTGCG
490	500	510	520	530	540
TCTGTCTTC	TAATAATGA	AGGACTACTT	GATAAATCAT	TTATGCTGAA	CGCATGTTAT
550	560	570	580	590	600
GATGTAATAA	GTGAAGGCAT	ACCTTTTCT	CCTTATATAT	GTGCAGGTAT	TGGTATTGAT
610	620	630	640	650	660
TTAGTATCCA	TGTTTGAAGC	TATAATCCT	AAAATTTCTT	ATCAAGGAAA	ATTAGGCTTA
670	680	690	700	710	720
AGTTACCTA	TAAGCCCAGA	AGCTTCTGTG	TTTATTGGTG	GACATTTCA	TAAGGTGATA
730	740	750	760	770	780
GGAAACGAAT	TTAGAGATAT	TCCTACTATG	ATACCTAGTG	AATCAGCGCT	TGCAGGAAAA
790	800	810	820	830	840
GGAAACTACC	CTGCAATAGT	AAACACTGGAC	GTGTTCTACT	TTGGCATAGA	ACTTGGAGGA
850	860	870	880	890	900
AGGTTTAACT	TCCAACTTG	A.....

Fig. 6A

10	20	30	40	50	60
MNCEKFFITT	ALTLMSFLP	GISLSDPVQD	DNISGNFYIS	GKYMP SASHF	GVFSAKEERN
70	80	90	100	110	120
TTVGVFGIEQ	DWDRCVISRT	TLSDFITVPN	YSFKYENNLF	SGFAGAIGYS	MDGPRIELEV
130	140	150	160	170	180
SYEAFDVKNQ	GNNYKNEAHR	YYALSHLLGT	ETQIDGAGSA	SFVLINEGLL	DKSFMLNACY
190	200	210	220	230	240
DVSEGIPFS	PYICAGIGID	LVSMFEAINP	KISYQGKLGL	SYPISPEASV	FIGGHFHVKVI
250	260	270	280	290	300
GNEFRDIPTM	IPSESALAGK	GNYPAIVTLD	VFYFGIELGG	RFNFQL

Fig. 6B

6/31

10	20	30	40	50	60
ATGAATTGCA	AAAAATTTT	TATAACAACT	GCATTAGTAT	CACTAATGTC	CTTTCTACCT
70	80	90	100	110	120
GGAATATCAT	TTTCTGATCC	AGTGCAAGGT	GACAATATTA	GTGGTAATTT	CTATGTTAGT
130	140	150	160	170	180
GGCAAGTATA	TGCCAAGTGC	TTCGCATT	GGCATGTTT	CTGCCAAAGA	AGAAAAAAT
190	200	210	220	230	240
CCTACTGTTG	CATTGTATGG	CTTAAAACAA	GATTGGGAAG	GGATTAGCTC	ATCAAGTCAC
250	260	270	280	290	300
AATGATAATC	ATTTCAATAA	CAAGGGTTAT	TCATTTAAAT	ATGAAATAA	CCCATTTTA
310	320	330	340	350	360
GGGTTTGCAG	GAGCTATTGG	TTATTCAATG	GGTGGTCCAA	GAGTAGAGTT	TGAAGTGTCC
370	380	390	400	410	420
TATGAAACAT	TTGACGTTAA	AAATCAGGGT	AATAACTATA	AAAATGATGC	TCACAGATAAC
430	440	450	460	470	480
TGTGCTTTAG	GTCAACAAGA	CAACAGCGGA	ATACCTAAAA	CTAGTAAATA	CGTACTGTTA
490	500	510	520	530	540
AAAAGCGAAG	GATTGCTTGA	CATATCATT	ATGCTAAATG	CATGCTATGA	TATAATAAAC
550	560	570	580	590	600
GAGAGCATAAC	CTTTGCTCTCC	TTACATATGT	GCAGGTGTTG	GTACTGATT	AATATCCATG
610	620	630	640	650	660
TTTGAAGCTA	CAAATCCTAA	AATTTCTTAC	CAAGGGAACT	TAGGTCTAAG	TTACTCTATA
670	680	690	700	710	720
AACCCAGAAG	CTTCTGTATT	TATTGGTGG	CATTTCTATA	AGGTGATAGG	AAACGAATT
730	740	750	760	770	780
AGGGACATTC	CTACTCTGAA	AGCATTGTT	ACGTCACTAG	CTACTCCAGA	TCTAGCAATA
790	800	810	820	830	840
GTAACACTAA	GTGTATGTCA	TTTTGGAATA	GAACTTGGAG	GAAGGTTAA	CTTCTAA...

Fig. 7A

10	20	30	40	50	60
MNCKKFFIIT	ALVSLMSFLP	GISFSDPVQG	DNIISGNFYVS	GKYMPASAHF	GMFSAKEEKN
70	80	90	100	110	120
PTVALYGLKQ	DWEGISSLSSH	NDNHENNNKGY	SFKYENNPF	GFAGAIGYSM	GGPRVFEFEVS
130	140	150	160	170	180
YETFDVKNQG	NNYKNDAHRY	CALGQODNSG	IPKTSKYVLL	KSEGLLDISF	MLNACYDIIN
190	200	210	220	230	240
ESIPLSPYIC	AGVGTDLISM	FEATNPKISY	QGKLGLSYSI	NPEASVFIGG	HFHKVIGNEF
250	260	270	280	290	300
RDIPTLKAFV	TSSATPDLAI	VTLSVCHFGI	ELGGRFNF...

Fig. 7B

7/31

10	20	30	40	50	60
ATGAATTGCA	AAAAATTTT	TATAACAACT	ACATTAGTAT	CGCTAATGTC	CTTCTTACCT
70	80	90	100	110	120
GGAATATCAT	TTTCTGATGC	AGTACAGAAC	GACAATGTTG	GTGGTAATTT	CTATATCAGT
130	140	150	160	170	180
GGGAAATATG	TACCAAGTGT	TTCACATTTT	GGCGTATTCT	CTGCTAAACA	GGAAAGAAAT
190	200	210	220	230	240
ACAACAAACCG	GAGTATTTGG	ATTAAAGCAA	GATTGGGATG	GCAGCACAAT	ATCTAAAAAT
250	260	270	280	290	300
TCTCCAGAAA	ATACATTTAA	CGTTCCAAAT	TATTCATTTA	AATATGAAAA	TAATCCATTT
310	320	330	340	350	360
CTAGGTTTG	CAGGAGCTGT	TGGTTATTTA	ATGAATGGTC	CAAGAATAGA	GTTAGAAATG
370	380	390	400	410	420
TCCTATGAAA	CATTTGATGT	AAAAAACCAG	GGTAATAACT	ATAAGAACGA	TGCTCACAAA
430	440	450	460	470	480
TATTATGCTT	TAACCCATAA	CAGTGGGGGA	AAGCTAAGCA	ATGCAGGTGA	TAAGTTGTT
490	500	510	520	530	540
TTTCTAAAAA	ATGAAGGACT	ACTTGATATA	TCACTTATGT	TGAATGCATG	CTATGATGTA
550	560	570	580	590	600
ATAAGTGAAG	GAATACCTTT	CTCTCCTTAC	ATATGTGCAG	GTGTTGGTAC	TGATTTAATA
610	620	630	640	650	660
TCCATGTTG	AAGCTATAAA	CCCTAAAATT	TCTTATCAAG	GAAAGTTAGG	TTTGAGTTAC
670	680	690	700	710	720
TCCATAAGCC	CAGAAGCTTC	TGTTTTGTT	GGTGGACATT	TTCATAAGGT	GATAGGGAAT
730	740	750	760	770	780
GAATTCAAGAG	ATATTCCCTGC	TATGATAACCC	AGTACCTCAA	CTCTCACAGG	TAATCACTTT
790	800	810	820	830	840
ACTATAGTAA	CACTAAGTGT	ATGCCACTTT	GGAGTGGAAC	TTGGAGGAAG	GTTTAACTTT
850	860	870	880	890	900
TAA.....

Fig. 8A

10	20	30	40	50	60
MNCKKKFIIT	TLVSLMSFLP	GISFSDAVQN	DNVGGNFYIS	GKYVPSVSHF	GVFSAKQERN
70	80	90	100	110	120
TTTGVFGLKQ	DWDGSTISKN	SPENTFNVPN	YSFKYENNPF	LGFAGAVGVL	MNGPRIELEM
130	140	150	160	170	180
SYETFDVKNQ	GNNYKNDARK	YYALTHNSGG	KLSNAGDKFV	FLKNEGLLDI	SIMLNACYDV
190	200	210	220	230	240
ISEGIPFSPY	ICAGVGTDLI	SMFEAINPKI	SYQGKLGLSY	SISPEASVVF	GGHFKVIGN
250	260	270	280	290	300
EFRDIPAMIP	STSTLTGNHF	TIVTLSVCHF	GVELGGRFNF

Fig. 8B

10	20	30	40	50	60
ATGGAAAATC	TCATGAATAA	GAAAAACAAA	TTCTTTACAA	TAAGTACAGC	AATGGTATGC
70	80	90	100	110	120
TTATTGTTAT	TACCTGGTAT	ATCATTTCA	GAACTATAA	ACAACAGTGC	TAAAAAACAG
130	140	150	160	170	180
CCTGGGTTAT	ATATCAGTGG	GCAGTACAAA	CCTAGTGTTC	CAGTTTTAG	TAATTTTCA
190	200	210	220	230	240
GTAAAAGAAA	CTAATGTTCC	CACAAAGCAG	TTAATAGCAC	TTAAAAAAGA	CATTAATTCT
250	260	270	280	290	300
GTTGCAGTTG	GTAGTAATGC	TACTACAGGT	ATTAGCAATC	CAGGTAATT	CACAATTCC
310	320	330	340	350	360
TATACTGCAG	AATTTCAAGA	TAATGTTGCC	AATTTCAATG	GGGCTGTTGG	TTACTCTTTT
370	380	390	400	410	420
CCTGATAGTC	TAAGAATTGA	AATAGAGGGA	TTTCATGAAA	AATTTGATGT	CAAAAACCC
430	440	450	460	470	480
GGAGGTTACA	CACAAGTAAA	AGATGCGTAC	CGTTATTTG	CACTAGCACG	TGATTTAAA
490	500	510	520	530	540
GATGGCTTCT	TTGAACCTAA	AGCGGAAGAT	ACAGGTGTTC	ATCATACTGT	TATGAAAAAT
550	560	570	580	590	600
GATGGATTAT	CTATTTATC	TACTATGGTT	AACGTCTGTT	ACGATTTTC	TGTAGATGAA
610	620	630	640	650	660
TTACCAAGTCT	TACCTTATAT	ATGTGCAGGT	ATGGGTATAA	ACGCCATAGA	ATTCTTCGAC
670	680	690	700	710	720
GCTTTACATG	AAAAATTTC	TTACCAAGGC	AAACTAGGTA	TTAGCTATCA	ACTATTTACT
730	740	750	760	770	780
AAAGTAAATT	TATTCCTTGA	TGGGTATTAC	CATCAAGTAA	TAGGCAATCA	ATTCAAAAC
790	800	810	820	830	840
TTAAACGTAA	ACCATGTTA	CACACTAAA	GAATCTCCTA	AAGTCACATC	TGCAGTAGCT
850	860	870	880	890	900
ACACTTGACA	TTGCATACTT	TGGTGGCGAA	GTTGGAATAA	GATTACATT	TTAA.....

Fig. 9A

10	20	30	40	50	60
MVCLLILPGI	SFSETINNSA	KKQPGLYISG	QYKPSVSVFS	NFSVKEINVP	TKQLIAALKRD
70	80	90	100	110	120
INSVAVGSNA	TTGISNPNGNF	TIPYTAEFQD	NVANFNGAVG	YSFPDSLRIE	IEGFHEKFDV
130	140	150	160	170	180
KNPGGYTQVK	DAYRYFALAR	DLKDGFEPK	AEDTGVYHTV	MKNDGLSILS	TMVNVCYDFP
190	200	210	220	230	240
VDELPVLPYI	CAGMGINAIE	FFDALHVKFA	YQGKLGISYQ	LFTKVNLFID	GYYHQVIGNQ
250	260	270	280	290	300
FKNLNVNVHVY	TLKESPKVTS	AVATLDIAYF	GGEVGIRFTF

Fig. 9B

9/31

10	20	30	40	50	60
ATGATATATA	AAGAAAAACT	TAATAGAGTG	GGAGAAATATA	TCTTAGCATA	TTTATCATT
70	80	90	100	110	120
ATTCTTTCTA	CTTATATCTT	TCTAGTGCTG	GTAAATATTA	TTAGATATAA	CAGCCTTGCT
130	140	150	160	170	180
ATATGTGTTA	TCAGTCTACT	AAGAACTAAT	ATCTTTAACG	TTAGCACAAA	AAAATTAATA
190	200	210	220	230	240
AAAGATAAAAT	GTCGTGATAC	TAAGTTAGT	AACATGAATT	GTTATTTGTA	CGGTAAACCG
250	260	270	280	290	300
TTAAATTTAC	AAATTTTTA	TGGAATATTT	TCCTTTATTA	GAAACTTTCA	AAATAACACA
310	320	330	340	350	360
CTAATAATTC	CTAATGATAG	TAAATGCGGC	TTCTATACCA	CGTTATGGGA	TAATCCAGCA
370	380	390	400	410	420
CTACATTATA	CATATACACT	TACTGGCACT	GAGTACCGTA	ATTTTTTGTA	CATTCTATAT
430	440	450	460	470	480
GAAAACATTA	TCTGTCAATG	TAAATTACTT	ATTAACTATA	ACCGTTCTGT	ATTAACCAA
490	500	510	520	530	540
CATAATAAAA	ATACTCTCGT	AATAATACCA	ATACCTAATG	CTAGAGAGTT	CAGTAATGAA
550	560	570	580	590	600
ATTCGAGTAA	GGAATATATC	AATAAATAAG	GAAAGTTCTT	ATGAGTGCTA	A.....

Fig. 10A

10	20	30	40	50	60
MIYKEKLTRV	GEYILAYLSF	ILSTYIFLVL	VNIIRYNSLA	ICVISLLRTN	IFNVSTKKLI
70	80	90	100	110	120
KDKCRDTKFS	NMNCYLYGKP	LNLQIFYGIF	SFIRNFQNNT	LIIPNDSKCG	FYTTLWDNPA
130	140	150	160	170	180
LHYTYTLTGS	EYRNFFDILY	ENIICQCKLL	INYNRSQLNQ	HNKNTLVIIP	IPNAREFSNE
190	200	210	220	230	240
IRVRNISINK	ESSYEC.....

Fig. 10B

10/31

10	20	30	40	50	60
ATGAATAAAA	AAAACAAGTT	TATTATAGCT	ACAGCATTGG	TATATTACT	GTCATTACCT
70	80	90	100	110	120
AGTGTATCGT	TTTCAGAGGT	TACAAACAGC	AGTATTAAAA	AACACTCTGG	GTTATATATT
130	140	150	160	170	180
AGTGGACAAT	ACAAACCAAG	TGTTTCTGTT	TTTAGTAGTT	TCTCAATTAA	AGAAAACAAAC
190	200	210	220	230	240
ACTATCACAA	AAAATCTTAT	AGCGTTAAAA	AAAGATATTA	ACTCTCTTGA	AGTTAACGCC
250	260	270	280	290	300
GATGCTAGTC	AAGGTATTAG	TCATCCAGGA	AATTTTACTA	TACCTTATAT	AGCAGCATT
310	320	330	340	350	360
GAAGATAATG	CTTTTAATTT	CAACGGTGCT	ATTGGTTACA	TTACTGAAGG	TCTAAGGATT
370	380	390	400	410	420
GAAATAGAAG	GTTCCATGA	AGAATTTGAT	GCTGAAAACC	CTGGAGGTTA	TGGTCTAAAT
430	440	450	460	470	480
GATGCCTTTC	GGTACTTTGC	TTTAGCACGT	GATATGGAAA	GCAACAAGTT	CCTACCAAAA
490	500	510	520	530	540
GCACAAAGCT	CAC.....				

Fig. 11A

10	20	30	40	50	60
MNKKNKFIIA	TALVYLLSLP	SVSFSEVTNS	SIKKHSGLYI	SGQYKPSVSV	FSSFSIKETN
70	80	90	100	110	120
TITKNLIALK	KDINSLEVNA	DASQGISHPG	NFTIPIYIAAF	EDNAFNFNGA	IGYITEGLRI
130	140	150	160	170	180
EIEGSYEEFD	AENPGGYGLN	DAFRYFALAR	DMESNKFLPK	AQSS.....	

Fig. 11B

10	20	30	40	50	60
70	80	90	100	110	120
130	140	150	160	170	180
190	200	210	220	230	240
250	260	270	280	290	300
310	320	330	340	350	360
370	380	390	400	410	420
GGAAGTGAAA TTGGAATAAG GTTTATGTTT AACTAA.....					

Fig. 12A

10	20	30	40	50	60
70	80	90	100	110	120
130	140	150	160	170	180
GSEIGIREMF N.....					

Fig. 12B

12/31

10	20	30	40	50	60
ATGACAAAGA	AATTAAATTT	TGTAAATGTT	ATATTAACAT	TTTTGTTATT	TCTTTCCCA
70	80	90	100	110	120
CTTAAGTCAT	TTACAAACATA	TGCAAAATAAT	AACACAATCA	CTCAAAAAGT	TGGATTGTAC
130	140	150	160	170	180
ATAAGTGGTC	AATATAAGCC	AAGTATTCCCT	CATTTCAAGA	ATTTTCAGT	AGAAGAAAAT
190	200	210	220	230	240
GACAAAGTAG	TAGATTGAT	AGGTCTTACA	ACTGATGTTA	CATATATCAC	AGAACATATA
250	260	270	280	290	300
TTACGAGATA	ATACAAAATT	CAACACTCAT	TATATTGCAA	AGTTCAAGAA	CAATTTATA
310	320	330	340	350	360
AATTCAGCA	GTGCAATTGG	TTATTATTCT	GGGCAAGGAC	CAAGGTTAGA	AATAGAAAGC
370	380	390	400	410	420
TCTTATGGGG	ATTTGATGT	TGTAAATTAT	AAAAATTATG	CAGTACAAGA	TGTTAATAGA
430	440	450	460	470	480
TATTTTGCTT	TAGTACGTGA	AAAAAATGGT	TCAAATTCT	CTCCAAAACC	ACATGAAACT
490	500	510	520	530	540
AGTCAACCCCT	CTGACAGTAA	TCCTAAAAAG	TCTTTTATA	CTTTAATGAA	GAATAATGGG
550	560	570	580	590	600
GTATTTGTTG	CATCAGTAAT	AATCAACGGT	TGTTATGATT	TTTCTTTAA	TAACACAAACA
610	620	630	640	650	660
ATATCACCTT	ACGTATGTAT	AGGAGTTGGA	GGAGATTTA	TAGAGTTTT	TGAAGTAATG
670	680	690	700	710	720
CATATCAAGT	TTGCTTGCCA	AAGTAAGGTT	GGTATTAGCT	ATCCAATATC	TCCCTCTATT
730	740	750	760	770	780
ACTATTTTG	CTGATGCACA	TTATCACAAG	GTCATAAATA	ATAAATTAA	CAACCTACAT
790	800	810	820	830	840
GTAAAGTATT	CATATGAACT	TAAAAACTCA	CCTACCATT	CCTCTGCAAC	AGCCAAACTA
850	860	870	880	890	900
AACATTGAAT	ATTTGGTGG	TGAAGTTGGG	ATGAGATTAA	TATTTAA

Fig. 13A

10	20	30	40	50	60
MTKKFNFVN	ILTFILLFLFP	LKSFTTYANN	NTITQKVGLY	ISGQYKPSIP	HFKNFSVEEN
70	80	90	100	110	120
DKVVVDLIGLT	TDVTYITEHI	LRONTKFNT	YIAKFNNFI	NFSSAIGYYS	GQGPRLEIES
130	140	150	160	170	180
SYGDFDVVNY	KNYAVQDVNR	YFALVREKNG	SNFSPKPHET	SQPSDSNPKK	SFYTLMKNNNG
190	200	210	220	230	240
VFVASVIING	CYDFSFNNTT	ISPYVCIGVG	GDFIEFFEV	HIKFACQSKV	GISYPISPSI
250	260	270	280	290	300
TIFADAHYHK	VINNKFNNLH	VKYSYELKNS	PTITSATAKL	NIEYFGGEVG	MRFIF.....

Fig. 13B

10	20	30	40	50	60
ATGAGCAAAA	AAAAGTTTAT	TACAATAGGA	ACAGTACTTG	CATCTCTATT	ATCATTCTTA
70	80	90	100	110	120
TCTATTGAAT	CCTTTTCAGC	TATAAATCAT	AATCATAACAG	GAAATAACAC	TAGTGGTATA
130	140	150	160	170	180
TATATTACAG	GGCAGTATAG	ACCAGGAGTA	TCCCATTAA	GCAATTCTC	AGTAAAAGAA
190	200	210	220	230	240
ACTAATGTTG	ATACAATACA	ACTAGTAGGA	TATAAAAAAA	GTGCGTCTTC	TATCGATCCT
250	260	270	280	290	300
AACACTTATT	CAAACTTCA	AGGTCCATAT	ACTGTTACAT	TTCAAGATAA	TGCTGCTAGT
310	320	330	340	350	360
TTCAGTGGAG	CAATTGGATA	TTCTTACCCC	GAAAGTCTAA	GACTTGAAC	TGAAGGTTCT
370	380	390	400	410	420
TACGAAAAAT	TTGATGTCAA	AGATCCTAAA	GACTACTCAG	CAAAGATGC	TTTTAGGTTT
430	440	450	460	470	480
TTTGCTCTAG	CACGTAATAC	GTCTACTACT	GTTCCCTGATG	CTCAAAAATA	TACAGTTATG
490	500	510	520	530	540
AAGAATAATG	GCTTATCTGT	TGCATCAATC	ATGATCAATG	GTTGTTATGA	TCTATCTTTT
550	560	570	580	590	600
AATAATTTAG	TCGTATCACC	TTATATATGT	GCAGGTATTG	GTGAAGATT	CATTGAATT
610	620	630	640	650	660
TTTGATACTT	TGCACATTAA	ACTTGCTTAT	CAAGGAAAAC	TAGGTATTAG	TTATTACTTC
670	680	690	700	710	720
TTTCCTAAGA	TTAATGTATT	TGCTGGGGGG	TACTATCATA	GAGTTATAGG	GAATAAATT
730	740	750	760	770	780
AAAAATTTAA	ATGTTAACCA	TGTTGTTACA	CTTGATGAAT	TTCCTAAAGC	AACTTCTGCA
790	800	810	820	830	840
GTAGCTACAC	TTAATGTTGC	TTATTTGGT	GGTGAAGCTG	GAGTAAAGTT	TACATTTAA
850	860	870	880	890	900
.....

Fig. 14A

10	20	30	40	50	60
MSKKKFITIG	TVLASLLSFL	SIESFSAINH	NKTGNNTSGI	YITGQYRPGV	SHFSNFSVKE
70	80	90	100	110	120
TNVDTIQLVG	YKKASSIDP	NTYSNFQGPY	TVTFQDNAAS	FSGAIGYSYP	ESLRLELEGS
130	140	150	160	170	180
YEKFVDVKDPK	DYSAKDAERF	FALARNTSTT	VPDAQKYTVM	KNNGLSVASI	MINGCYDLSE
190	200	210	220	230	240
NNLVVSPYIC	AGIGEDFIEF	FDTLHIKLAY	QGKLGISYYF	FPKINVEAGG	YHHRVIGNKE
250	260	270	280	290	300
KNLNVNHVVT	LDEFPKATSA	VATLNVAYFG	GEAGVKFTF

Fig. 14B

10	20	30	40	50	60
ATGAGTGCTA	AAAAAAAGCT	TTTTATAATA	GGGTCAAGTGT	TAGTATGTTT	AGTGTCACTAC
70	80	90	100	110	120
TTACCTACTA	AATCTTGTC	AAACTTAAAT	AATATTAAATA	ATAACACTAA	GTGCACTGGG
130	140	150	160	170	180
CTATATGTCA	GTGGACAATA	TAAACCTACT	GTTCCTCACT	TTAGTAATT	TTCACTTAA
190	200	210	220	230	240
GAAACTTATA	CTGACACTAA	AGAGTTATTA	GGACTAGCAA	AAGATATTAA	GTCTATTACA
250	260	270	280	290	300
GATATAACAA	CAAATAAAAA	ATTCAACATT	CCTTATAACA	CAAAATTCA	AGATAATGCT
310	320	330	340	350	360
GTTAGCTTCA	GTGCAGCTGT	TGGATATATT	TCCCAAGACA	GTCCAAGGGT	TGAGGTAGAA
370	380	390	400	410	420
TGGTCTTATG	AAGAATTGTA	CGTTAAAAAT	CCTGGTAATT	ACGTAGTAAG	TGAAGCCTTC
430	440	450	460	470	480
AGGTATATTG	CTTTAGCAAG	AGGAATTGAT	AATCTTCAAA	AATATCCTGA	AACAAATAAG
490	500	510	520	530	540
TATGTTGTTA	TAAAGAACAA	TGGCTTATCT	GTCGCATCCA	TTATAATCAA	TGGCTGTTAT
550	560	570	580	590	600
GATTTTTCTT	TAAACAATTT	AAAAGTATCA	CCTTACATAT	GCGTAGGGTT	TGGTGGGGAC
610	620	630	640	650	660
ATTATAGAAT	TTTTTAGTGC	TGTAAGTTT	AAATTTGCTT	ATCAAGGTAA	GGTAGGTATC
670	680	690	700	710	720
AGTTATCCAT	TATTCTCTAA	TATGATTATA	TTTGCTGACG	GATATTACCA	TAAGGTCAAA
730	740	750	760	770	780
GGAAATAAAAT	TTAACAAATTT	AAATGTTCAA	CACGTTGTTA	GTCTAACAG	TCATCCTAAAG
790	800	810	820	830	840
TCTACTTTTG	CAGTAGCTAC	TCTTAATGTT	GAGTATTTCG	GTAGTGAATT	TGGGTTAAAA
850	860	870	880	890	900
TTTATATTTT AA.....					

Fig. 15A

10	20	30	40	50	60
MSAKKKLFII	GSVILVCLVSY	LPTKSLSNLN	NINNNNTKCTG	LYVSGQYKPT	VSHFSNFSLK
70	80	90	100	110	120
ETYTDTKELL	GLAKDIKSIT	DITTNKFQDNA	VSFSAAVGYI	SQDSPRVEVE	
130	140	150	160	170	180
WSYEEFDVKN	PGNYVVSEAF	RYIALARGID	NLQKYPETNK	YVVIKNNGLS	VASIIINGCY
190	200	210	220	230	240
DFSLNNLKVS	PYICVGFGGD	IEFFSAVSE	KFAYQGKVGI	SYPLFSNMII	FADGYYHKVI
250	260	270	280	290	300
GNKFNNLNQ	HVVSLNSHPK	STFAVATLNV	EYFGSEFGLK	FIF.....	

Fig. 15B

10	20	30	40	50	60
ATGAGTAAAA	AAAATTTAT	TACAATAGGA	GCAACACTTA	TTCATATGTT	GTTACCTAAC
70	80	90	100	110	120
ATATCTTTTC	CAGAACTAT	TAACAATAAC	ACTGATAAAC	TTTCTGGGTT	ATATATAAGT
130	140	150	160	170	180
GGGCAATATA	AACCAGGGAT	TTCTCATTTC	AGCAAATTT	CAGTCAAAGA	AATCTATAAT
190	200	210	220	230	240
GATAACATTC	AACTAATTGG	GTAAAGACAC	AACGCAATTT	CTACTAGTAC	CCTTAATATT
250	260	270	280	290	300
AATACAGATT	TTAAATATCCC	CTATAAAGTA	ACATTTCAAA	ATAACATTAC	CAGCTTTAGT
310	320	330	340	350	360
GGAGCTATTG	GTTATTCTGA	TCCCACAGGG	GCAAGATTG	AGCTTGAAGG	TTCTTATGAA
370	380	390	400	410	420
GAATTTGATG	TGACAGATCC	TGGAGACTGC	TTAATAAAAG	ATACCTATAG	ATATTCGCT
430	440	450	460	470	480
TTAGCTAGAA	ACCCATCAGG	TTCTAGCCCT	ACCTCAAACA	ACTATACTGT	TATGAGAAAT
490	500	510	520	530	540
GATGGTGT	CCATTACTTC	TGTTATATTT	AATGGCTGTT	ATGACATCTT	TTAAAGGAT
550	560	570	580	590	600
TTAGAAGTAT	CACCTTATGT	ATGTGTTGGT	GTAGGTGGAG	ATTTTATAGA	ATTTTTGAC
610	620	630	640	650	660
GCATTACACA	TTAAATTAGC	ATACCAAGGC	AAGTTAGGTA	TCAATTATCA	CTTATCGACT
670	680	690	700	710	720
CAAGCAAGCG	TATTTATTGA	TGGATATTAT	CATAAGGTTA	TAGGAAATCA	ATTCAACAAT
730	740	750	760	770	780
CTAAATGTC	AACACGTGGC	TAGTACAGAT	TTTGGACCTG	TATACGCAGT	AGCCACACTT
790	800	810	820	830	840
AACATTGGTT ATTTTGGTGG TGAAATCGGA ATTAGACTTA CATTAA..					

Fig. 16A

10	20	30	40	50	60
MSKKNFITIG	ATLIHMLLPN	ISFPETINNN	TDKLSGLYIS	GQYKPGISHF	SKFSVKEIYN
70	80	90	100	110	120
DNIQLIGLRH	NAISTSTLNI	NTDENIPYKV	TFQNNNITSFS	GAIGYSDPTG	ARFELEGSYE
130	140	150	160	170	180
EFDVTDPGDC	LIKDTYRYFA	LARNPSGSSP	TSNNYTVMRN	DGVSITSVIF	NGCYDIFLKD
190	200	210	220	230	240
LEVSPYVCVG	VGGDFIEFFD	ALHIKLAYQG	KLGINYHLST	QASVFIGDYY	HKVIGNQFNN
250	260	270	280	290	300
INVQHVASTD	FGPVYAVATL	NIGYFGGEIG	IRLTF

Fig. 16B

10	20	30	40	50	60
ATGAATAATA	GAAAAAGTTT	TTTTATAATA	GGTGCATCAT	TACTAGCAAG	CTTATTATTTC
70	80	90	100	110	120
ACATCTGAGG	CCTCTTCTAC	AGGAAATGTA	AGTAACCATA	CTTATTITAA	ACCTAGGTTA
130	140	150	160	170	180
TATATCAGTG	GACAATATAG	ACCAGGAGTT	TCTCATTTTA	GCAAATTTTC	AGTCAAAGAA
190	200	210	220	230	240
ACCAACTACA	ATACTACTCA	ACTAGTTGGG	CTTAAAAGG	ACATCAGTGT	CATAGGGAAC
250	260	270	280	290	300
AGTAATATCA	CAACCTACAC	AAATTTCAAC	TTTCCTTACA	TTGCAGAATT	TCAAGACAAAT
310	320	330	340	350	360
GCCATAAGTT	TCAGTGGGGC	AATTGGATAC	TTGTATTCCG	AGAATTTCAG	AATTGAAGTA
370	380	390	400	410	420
GAGGCTTCTT	ATGAAGAATT	TGATGTTAAA	AATCCAGAAG	GATCTGCTAC	AGACGCATAC
430	440	450	460	470	480
AGGTATTTG	CACTAGCACG	TGCTATGGAT	GGCACTAATA	AATCTAGTCC	TGATGACACA
490	500	510	520	530	540
AGAAAATTCA	CTGTCATGAG	AAATGACGGG	TTATCAATT	CATCAGTAAT	GATAAATGGG
550	560	570	580	590	600
TGTTACAATT	TTACATTAGA	TGATATACCA	GTAGTACCGT	ATGTATGCGC	AGGAATAGGA
610	620	630	640	650	660
GGAGATTTCA	TAGAGTTTT	TAATGATT	CATGTTAAGT	TTGCTCATCA	AGGCAAGGTA
670	680	690	700	710	720
GGTATTAGTT	ATTCTATATC	CCCTGAAGTA	AGTTTATTTC	TTAACGGATA	TTACCATAAA
730	740	750	760	770	780
GTAACAGGTA	ACAGATTAA	AAACTTACAC	GTTCAACACG	TAAGTGATT	AAGTGACGCT
790	800	810	820	830	840
CCTAAGTTCA	CATCTGCAGT	TGCTACACTC	AATGTTGGGT	ACTTTGGTGG	CGAAATTGGGA
850	860	870	880	890	900
GTAAAGATTAA	TATTTAA..

Fig. 17A

10	20	30	40	50	60
MNNRKSFFII	GASILLASLLF	TSEASSTGNV	SNHTYFKPRL	YISGQYRPGV	SHFSKFSVKE
70	80	90	100	110	120
TNYNTTQLVG	LKKDISVIGN	SNITTYTNEN	FPYIAEFQDN	AISFSGAIGY	LYSENFRIEV
130	140	150	160	170	180
EASYEEFDVK	NPEGSATDAY	RYFALARAMD	GTNKSSPDDT	RKFTVMRNDG	LSISSVMING
190	200	210	220	230	240
CYNFTLDDIP	VVPYVCAGIG	GDFIEFFNDL	HVKFAHQGKV	GISYSISPEV	SLFLNGYYHK
250	260	270	280	290	300
VTGNRFKNLH	VQHVSDSLSDA	PKETSATVL	NVGYEGGEIG	VRFIF

Fig. 17B

10	20	30	40	50	60
TAGCAGCACT	AAAAAACAGT	TTGGGTTATA	TGTTAGTGGA	CAACACCAGC	CTAGTGTTTC
70	80	90	100	110	120
TATTTTTAGC	AATTCTCAAG	TAAAGGAAC	TAATTTTCCT	ACAAAGTATT	CTAGCAGCTT
130	140	150	160	170	180
CTTAAAAAAA	GACATTAATT	CTGTCGAATT	TGACGATAGT	GTTACTGCTG	GCATTAGTTA
190	200	210	220	230	240
CCCACCTTAAT	TTCAGTACTC	CTTATATAGC	TGTATTTCAA	GATAATATT	CTAATTTAA
250	260	270	280	290	300
TGGCGCTATT	GGGTACACTT	TTGTTGAAGG	CCCAAGAATT	GAAATAGAAG	GTTCTTATGA
310	320	330	340	350	360
AGAATTCCGAT	GTCAAAGACC	CTGGAAGATA	TACAGAAATA	CAAGATGCAT	ACCGTTACTT
370	380	390	400	410	420
TGCTTTAGCA	CGTGATATAG	ACTCTATTCC	TACTAGCCCA	AAAAATAGAA	CTTCACATGA
430	440	450	460	470	480
TGGCAACAGT	TCATATAAGG	TATACCACAC	TGTAATGAAA	AATGAAGGAC	TATCTATAAT
490	500	510	520	530	540
ATCCATTATG	GTCAATGGCT	GCTATGATTT	TTCTTCAGAT	AATTTATCAA	TATTACCTTA
550	560	570	580	590	600
TGTATGTGGT	GGTATAGGTG	TAATGCTAT	AGAGTTTTTC	GATGCATTAC	ATGTTAAATT
610	620	630	640	650	660
CGCGTGTCAAG	GGTAAATTAG	GTATTACTTA	TCCATTATCT	TCCAACGTTA	GTTTATTTGC
670	680	690	700	710	720
TGGTGGATAT	TATCACCAAG	TAATGGCAA	CCAATTAAA	AATCTAAATG	TTCACATGT
730	740	750	760	770	780
AGCTGAACCTT	AATGACGCAC	CCAAAGTTAC	ATCTGCAGTA	GCTACACTTG	ACATTGGGTA
790	800	810	820	830	840
TTTGGTGGT GAAATTGGAG CAAGGCTTAT ATTTTAA.....					

Fig. 18A

10	20	30	40	50	60
SSTKKQFGLY	VSGQHQPSVS	IFSNFSVKET	NFPTKYSSSF	LKKDINSVEF	DDSVTAGISY
70	80	90	100	110	120
PLNFSTPYIA	VFQDNISNPN	GAIGYTFVEG	PRIEIEGSYE	EFDVKDPGRY	TEIQDAYRYF
130	140	150	160	170	180
ALARIDIDSIP	TSPKNRTSHD	GNSSYKVYHT	VMKNEGLSII	SIMVNGCYDF	SSDNLSILPY
190	200	210	220	230	240
VCGGIGVNAI	EFFDALHVKF	ACQGKLGITY	PLSSNVSLFA	GGYYHQVMGN	QFRNLNVQHV
250	260	270	280	290	300
AELNDAPKVT SAVATLDIGY FGGEIGARLI F.....					

Fig. 18B

10	20	30	40	50	60
ATGAATTGCA	AAAGATTTT	CATAGCAAGT	GCATTGATAT	CACTAATGTC	TTTCTTACCT
70	80	90	100	110	120
AGCGTATCTT	TTTCTGAATC	AATACATGAA	GATAATATAA	ATGGTAACTT	TTACATTAGT
130	140	150	160	170	180
GCAAAGTATA	TGCCAAGTGC	CTCACACTT	GGCGTATTT	CAGTTAAAGA	AGAGAAAAAC
190	200	210	220	230	240
ACAACAACTG	GAGTTTCGG	ATTAACACAA	GATTGGGACG	GAGCAACAAT	AAAGGATGCA
250	260	270	280	290	300
AGCAGCAGCC	ACACAATAGA	CCCAAGTACA	ATATTCTCCA	TTTCAAATT	TTCATTTAAA
310	320	330	340	350	360
TATGAAAACA	ATCCATTTC	AGGGTTTGCA	GGAGCTATTG	GCTACTCAAT	GGGTGGTCCA
370	380	390	400	410	420
AGGGTAGAGT	TTGAAGTGT	TTACGAAATA	TTTGATGTAA	AAAACCAAGG	TAACAGTTAC
430	440	450	460	470	480
AAGAACGATG	CTCACAAATA	TTGCGCTTT	TCAAGACACA	CCGGAGGTAT	GCCACAAGCC
490	500	510	520	530	540
GGTCATCAA	ATAAATTGT	CTTCCTAAA	AATGAAGGAT	TACTTGACAT	ATCACTTATG
550	560	570	580	590	600
ATAAACGCAT	GTTATGATAT	AACAATCGAC	AGCATGCCAT	TTTCTCCATA	TATATGTGCA
610	620	630	640	650	660
GGTATTGGTA	GTGACTTAGT	TTCGATGTT	GAAACTACAA	ATCCTAAAAT	TTCTTATCAA
670	680	690	700	710	720
GGAAAATTAG	GTGTAAGTTA	CTCCATAAGC	CCAGAAGCAT	CTGTTTTGT	TGGAGGACAC
730	740	750	760	770	780
TTTCACAGAG	TTATAGGTAA	TGAATTAAA	GACATTCCGT	CAATAACTCC	TGCTGGAGCA
790	800	810	820	830	840
ACAGAAATTA	AAGGCACACA	GTTCACAACA	GTAACATTAA	ACATATGCCA	CTTCGGACTA
850	860	870	880	890	900
GAGCTTGGAG GCAGGTTAC TTTTAA...					

Fig. 19A

10	20	30	40	50	60
MNCKRFFIAS	ALISLMSFLP	SVSFSESIHE	DNINGNFYIS	AKYMPSASHF	GVFSVKEEKN
70	80	90	100	110	120
TTTGVFGLQ	DWDGATIKDA	SSSHTIDPST	IFSISNYSFK	YENNPFGLFA	GAIGYSMGGP
130	140	150	160	170	180
RVEFEVSYEI	FDVKNQGNSY	KNDAHKYCAL	SRHTGGMPQA	GHQNKFVFLK	NEGLLDISLM
190	200	210	220	230	240
INACYDITID	SMPFSPYICA	GIGSDLVSMF	ETTNPKISYQ	GKLGVSYYSIS	PEASVEVGHH
250	260	270	280	290	300
FHRVIGNEFK DIPAITPAGA TEIKGTQFTT VTINICHFGL ELGGRFTF..					

Fig. 19B

10	20	30	40	50	60
ATGAAATATA	AAAAACTTT	TACAGTAAC	GCATTAGTAT	TATTAACCTC	CTTTACACAT
70	80	90	100	110	120
TTTATACCTT	TTTATAGTCC	AGCACGTGCC	AGTACAATT	ACAACTTCTA	CATTAGTGGA
130	140	150	160	170	180
AAATATATGC	CAACAGCGTC	ACATTTGGA	ATTTTTTCA	CTAAAGAAGA	ACAAAGTTTT
190	200	210	220	230	240
ACTAAGGTAT	TAGTTGGGTT	AGATCAACGA	TTATCACATA	ATATTATAAA	CAATAATGAT
250	260	270	280	290	300
ACAGCAAAGA	GTCTTAAGGT	TCAAAATTAT	TCATTTAAAT	ACAAAATAA	CCCATTTCTA
310	320	330	340	350	360
GGATTTGCAA	GAGCTATTGG	TTATTCAATA	GGCAATTCAA	GAATAGAACT	AGAAGTATCA
370	380	390	400	410	420
CATGAAATAT	TTGATACTAA	AAACCCAGGA	AAACATTATT	TAAATGACTC	TCACAAATAT
430	440	450	460	470	480
TGCGCTTAT	CTCATGGAAG	TCACATATGC	AGTGATGGAA	ATAGCGGAGA	TTGGTACACT
490	500	510	520	530	540
GCAAAAAC	ATAAGTTGT	ACTTCTGAAA	AATGAAGGTT	TACTTGACGT	CTCATTTATG
550	560	570	580	590	600
TTAACCGCAT	GTTATGACAT	AACAAC	TGAA	AAAATGCCTT	TTTCACCTTA
610	620	630	640	650	660
GGTATTGGTA	CTGATCTCAT	ATCTATGTTT	GAGACAACAC	AAAACAAAAT	ATCTTATCAA
670	680	690	700	710	720
GGAAAGTTAG	GTTTAAACTA	TACTATAAAC	TCAAGAGTTT	CTGTTTTG	AGGTGGGCAC
730	740	750	760	770	780
TTTCATAAAG	TAATAGGTAA	TGAATTAAA	GGTATTCTA	CTCTATTACC	TGATGGATCA
790	800	810	820	830	840
AACATTAAAG	TACAACAGTC	TGCAACAGTA	ACATTAGATG	TGTGCCATT	CGGGTTAGAG
850	860	870	880	890	900
ATTGGAAGTA	GATTTTCTT	TTAA

Fig. 20A

10	20	30	40	50	60
MKYKKTFTVT	ALVLLTSFTH	PIPEYSPARA	STIHNFYISG	KYMPТАSHFG	IFSAKEEQSF
70	80	90	100	110	120
TKVLVGLDQR	LSHNIINNN	TAKS	SLKVQNY	SFKYKNMPEL	GEARAIGYSI
130	140	150	160	170	180
HEIFDTKMPG	NNYLNDSHKY	CALSHGSHIC	SDGNSGDWYT	AKTDKFVLLK	NEGLLDVSEM
190	200	210	220	230	240
LNACYDITTE	KMPPFS	SPYICA	GIGTDLISME	ETTONKISYQ	GKIGLNYTTIN
250	260	270	280	290	300
FKVIGNNEFK	GIPT	LLPDGS	NIKVQQSATV	TLDVCHFGL	IGSRFFF

Fig. 20B

10	20	30	40	50	60
ATGTTTTATA	CTAATATATA	TATTCTGGCT	TGTATTTACT	TTGCACCTTCC	ACTATTGTTA
70	80	90	100	110	120
ATTTATTTTC	ACTATTTAG	GTGTAATATG	AATTGCAAAA	AAATTCTTAT	AACAACGTGCA
130	140	150	160	170	180
TTAATATCAT	TAATGTACTC	TATTCCAAGC	ATATCTTTT	CTGATACTAT	ACAAGATGGT
190	200	210	220	230	240
AACATGGGTG	GTAACCTCTA	TATTAGTGGGA	AAGTATGTAC	CAAGTGTCTC	ACATTTGGT
250	260	270	280	290	300
AGCTTCTCAG	CTAAAGAAGA	AAGCAAATCA	ACTGTTGGAG	TTTTGGATT	AAAACATGAT
310	320	330	340	350	360
TGGGATGGAA	GTCCAATACT	TAAGAATAAA	CACGCTGACT	TTACTGTTCC	AAACTATTG
370	380	390	400	410	420
TTCAGATAACG	AGAACAAATCC	ATTTCTAGGG	TTTGCAGGAG	CTATCGGTTA	CTCAATGGGT
430	440	450	460	470	480
GGCCCAAGAA	TAGAATTCGA	AATATCTTAT	GAAGCATTG	ACGTAAAAAG	TCCTAATATC
490	500	510	520	530	540
AATTATCAA	ATGACGCGCA	CAGGTACTGC	GCTCTATCTC	ATCACACATC	GGCAGCCATG
550	560	570	580	590	600
GAAGCTGATA	AATTGTCTT	CTTAAAAAAC	GAAGGGTTAA	TTGACATATC	ACTTGCAATA
610	620	630	640	650	660
AATGCATGTT	ATGATATAAT	AAATGACAAA	GTACCTGTTT	CTCCTTATAT	ATGCGCAGGT
670	680	690	700	710	720
ATTGGTACTG	ATTTGATTTC	TATGTTGAA	GCTACAAGTC	CTAAAATTTTC	CTACCRAGGA
730	740	750	760	770	780
AAACTGGGCA	TTAGTTACTC	TATTAATCCG	GAAACCTCTG	TTTCATCGG	TGGGCATTTC
790	800	810	820	830	840
CACAGGATCA	TAGGTAATGA	GTTTAGAGAT	ATTCCCTGCAA	TAGTRACCTAG	TAACTCAACT
850	860	870	880	890	900
ACAATAAGTG	GACCACAATT	TGCAACAGTA	ACACTTAAATG	TGTGTCACTT	TGGTTTAGAA
910	920	930	940	950	960
CTTGGAGGAA	GATTTAACTT	CTAA

Fig. 21A

10	20	30	40	50	60
MFYTNIYILA	CIYFALPLLL	IYFHYFRCNM	NCKKILITTA	LISIMYSIPS	ISFSDTIQDG
70	80	90	100	110	120
NMGGNFYISG	KYVPSVSHEG	SFSAKEEKS	TVGVFGLKHD	WDGSPILKNK	HADFTVPNYS
130	140	150	160	170	180
FRYENNPFGL	FAGAIGYSMG	GPRIEFELSY	EAEDVKSPNI	NYQNDAHRYC	ALSHHTSAAM
190	200	210	220	230	240
EADKFVFLKN	EGLIDISLAI	NACYDIINDK	VPVSPYICAG	IGTDLISMEE	ATSPKISYQG
250	260	270	280	290	300
KLGISYSINP	ETSVFIGGHF	HRIIGMEFRD	IPAIVPSNST	TISGPQFATV	TLMVCHFGLE
310	320	330	340	350	360
LGGRNF...

Fig. 21B

10	20	30	40	50	60
ATGAATTGCA	AAAAAATTCT	TATAACAACT	GCATTAATGT	CATTAATGTA	CTATGCTCCA
70	80	90	100	110	120
AGCATACTT	TTTCTGATAC	TATACAAGAC	GATAACACTG	GTAGCTTCTA	CATCAGTGGA
130	140	150	160	170	180
AAATATGTAC	CAAGTGTTC	ACATTTGGT	GTTTCTCAG	CTAAAGAAGA	AAGAAACTCA
190	200	210	220	230	240
ACTGTTGGAG	TTTTGGATT	AAAACATGAT	TGGAATGGAG	GTACAATATC	TAACTCTTCT
250	260	270	280	290	300
CCAGAAAATA	TATTCACAGT	TCAAAATTAT	TCGTTAAAT	ACGAAAACAA	CCCATTCTTA
310	320	330	340	350	360
GGGTTTGCAG	GAGCTATTGG	TTATTCAATG	GGTGGCCCAA	GAATAGAACT	TGAAGTTCTG
370	380	390	400	410	420
TACGAGACAT	TCGATGTGAA	AAATCAGAAC	AATAATTATA	AGAACGGCGC	ACACAGATAAC
430	440	450	460	470	480
TGTGCTTTAT	CTCATCATAG	TTCAGCAACA	AACATGTCCT	CCGCAAGTAA	CAAATTTGTT
490	500	510	520	530	540
TTCTTAAAAA	ATGAAGGGTT	AATTGACTTA	TCATTATGA	TAAATGCATG	CTATGACATA
550	560	570	580	590	600
ATAATTGAAG	GAATGCCTTT	TTCACCTTAT	ATTTGTGCAG	GTGTTGGTAC	TGATGTTGTT
610	620	630	640	650	660
TCCATGTTTG	AAGCTATAAA	TCCTAAAATT	TCTTACCAAG	GAAAACTAGG	ATTAGGTTAT
670	680	690	700	710	720
AGTATAAGTT	CAGAACCTC	TGTTTTTATC	GGTGGACACT	TTCACAGAGT	CATAGGTAAT
730	740	750	760	770	780
GAATTTAGAG	ACATCCCTGC	TATGGTTCCCT	AGTGGATCAA	ATCTTCCAGA	AAACCAATT
790	800	810	820	830	840
GCAATAGTAA	CACTAAATGT	GTGTCACTTT	GGTTAGAAC	TTGGAGGAAG	ATTTAACTTC
850	860	870	880	890	900
TGA.....					

Fig. 22A

10	20	30	40	50	60
MNCKKILITT	ALMSLMYYAP	SISFSDTIQQ	DNTGSFYISG	KYVPSVSHFG	VFSAKEERNS
70	80	90	100	110	120
TVGVFGLKHD	WNGGTISNSS	PENIFTVQNY	SFKYENNPFL	GFAGAIGYSM	GGPRIELEV
130	140	150	160	170	180
YETFDVKNQN	NNYKNGAHRY	CALSHHSSAT	NMSSASNKFV	FLKNEGLIDL	SFMINACYDI
190	200	210	220	230	240
IIEGMPFSPY	ICAGVGTDVV	SMFEAINPKI	SYQGKLGLGY	SISSEASVFI	GGHFHRVIGN
250	260	270	280	290	300
EERDIPAMVP	SGSNLPEPQF	AIVTLMVCHF	GLELGGRFNF

Fig. 22B

10	20	30	40	50	60
ATGAATTGTA	AAAAAGTTT	CACAATAAGT	GCATTGATAT	CATCCATATA	CTTCCTACCT
70	80	90	100	110	120
AATGTCTCAT	ACTCTAACCC	AGTATATGGT	AACAGTATGT	ATGGTAATTT	TTACATATCA
130	140	150	160	170	180
GGAAAGTACA	TGCCAAGTGT	TCCTCATT	GGAATTTTT	CAGCTGAAGA	AGAGAAAAAA
190	200	210	220	230	240
AAGACAACTG	TAGTATATGG	CTTAAAAGGA	AAACTGGCAG	GAGATGCAAT	ATCTAGTC
250	260	270	280	290	300
AGTCCAGATG	ATAATTTTAC	CATTCGAAT	TACTCATTCA	AGTATGCAAG	CAACAAGTTT
310	320	330	340	350	360
TTAGGGTTTG	CAGTAGCTAT	TGGTTACTCG	ATAGGCAGTC	CAAGAATAGA	AGTTGAGATG
370	380	390	400	410	420
TCTTATGAAG	CATTTGATGT	GAAAAATCCA	GGTGATAATT	ACAAAAACGG	TGCTTACAGG
430	440	450	460	470	480
TATTGTGCTT	TATCTCATCA	AGATGATGCG	GATGATGACA	TGACTAGTGC	AACTGACAAA
490	500	510	520	530	540
TTTGTATATT	TAATTAATGA	AGGATTACTT	AACATATCAT	TTATGACAAA	CATATGTTAT
550	560	570	580	590	600
GAAACAGCAA	GCAAAAATAT	ACCTCTCTCT	CCTTACATAT	GTGCAGGTAT	TGGTACTGAT
610	620	630	640	650	660
TTAACATCACA	TGTTTGAAAC	TACACATCCT	AAAATTTCTT	ATCAAGGAAA	GCTAGGGTTG
670	680	690	700	710	720
GCCTACTTCG	TAAGTGCAGA	GTCTTCGGTT	TCTTTGGTA	TATATTTC	AAAATTATA
730	740	750	760	770	780
AATAATAAGT	TTAAAAATGT	TCCAGCCATG	GTACCTATTA	ACTCAGACGA	GATAGTAGGA
790	800	810	820	830	840
CCACAGTTG	CAACAGTAAC	ATTAAATGTA	TGCTACTTTG	GATTAGAACT	TGGATGTAGG
850	860	870	880	890	900
TTCAACTTCT	AA.....

Fig. 23A

10	20	30	40	50	60
MNCKKVFTIS	ALISSIYFLP	NVSYSNPVY	NSMYGNFYIS	GKYMPSPVHF	GIFSAEEEKK
70	80	90	100	110	120
KTIVVYGLKG	KLAGDAISSQ	SPDDNFTIRN	YSEKYASNKF	LGFAVAIGYS	IGSPRIEVEM
130	140	150	160	170	180
SYEAFDVKNP	GDNYKNGAYR	YCALSHQDDA	DDDMTSATDR	FVYLINEGLL	NISFMNTNICY
190	200	210	220	230	240
ETASKNIP	LSYICAGIGTD	LIHMFETTHP	KISYQGKLGL	AYFVSAESSV	SFGIYFHKII
250	260	270	280	290	300
NNKFKNPAM	VPINSDEIVG	PQFATVTLNV	CYFGLELGCR	FNF.....

Fig. 23B

10	20	30	40	50	60
ATGAACTGTA	AAAAATTCT	TATAACA	ACTAC	ACATTGGT	ACTAAACA
70	80	90	100	110	120
GGCATATCTT	TCTCCAAACC	AATACATGAA	AACAATACTA	CAGGAAACTT	TTACATTATT
130	140	150	160	170	180
GGAAAATATG	TACCAAGTAT	TTCACATT	GGGAAC	CAGCTAAAGA	AGAAAAAAAC
190	200	210	220	230	240
ACAACTACTG	GAATTTGG	ATTAAGA	TCATGGACTG	GTGGTATCAT	CCTTGATA
250	260	270	280	290	300
GAACATGCAG	CTTTAATAT	CCCAAATTAT	TCATTAA	ATGAAAATAA	TCCATT
310	320	330	340	350	360
GGATTTGCAG	GGGTAATTGG	CTATTCAATA	GGTAGTCCAA	GAATAGAATT	TGAAGTATCA
370	380	390	400	410	420
TACGAGACAT	TCGATGTACA	AAATCCAGGA	GATAAGTTA	ACAATGATGC	ACATAAGTAT
430	440	450	460	470	480
TGTGCTTAT	CCAATGATT	CAGTAAAACA	ATGAAAAGT	GTAAATT	TTTCTCA
490	500	510	520	530	540
AATGAAGGAT	TAAGTGACAT	ATCACTCATG	TTAAATGTAT	GT	TATGATAT
550	560	570	580	590	600
AGAATGCCCT	TTTCACCTTA	CATATGTGCA	GGCATTGGTA	CTGACTTA	ATTCA
610	620	630	640	650	660
GACGCTATAA	ACCATAAAGC	TGCTTATCAA	GGAAAATTAG	GT	TTTAATTA
670	680	690	700	710	720
CCAGAAGCTA	ACATTCTAT	GGGTGTGCA	TTTCACAAAG	TAACAAACAA	CGAGTTAGA
730	740	750	760	770	780
GTTCCTGTT	TATTAAC	TCGAGGACTC	GCTCCAGATA	ATCTATT	TGC
790	800	810	820	830	840
TTGAGTATAT	GTCATT	GGGTACAGGG	TCAGTT	TTA	A.....

Fig. 24A

10	20	30	40	50	60
MNCKKELITT	TLVSLTILLP	GISFSKPIHE	NNTTGNFYII	GKYVPSISHF	GNFSAKEEKN
70.	80	90	100	110	120
TTTGIFGLKE	SWTGGIILDK	EHAAFNIPNY	SFKYENNPL	GFAGVIGYSI	GSPRIEEVS
130	140	150	160	170	180
YETFDVQNP	DKFNNDAHKY	CALSNDSSKT	MKSGKEVFLK	NEGLSDISIM	LNVCYDIINK
190	200	210	220	230	240
RMPFSPYICA	GIGTDLIFMF	DAINHKAAYQ	GKLGFNYPIS	PEANISMGVH	EHKVTNNEFR
250	260	270	280	290	300
VPVLLTAGGL	APDNLFAIVK	LSICHFGLEF	GYRVSF

Fig. 24B

24/31

10	20	30	40	50	60
ATGAATAATA	AACTCAAATT	TACTATAATA	AACACAGTAT	TAGTATGCTT	ATTGTCATTA
70	80	90	100	110	120
CCTAATATAT	CTTCCTCAAA	GGCCATAAAC	AATAACGCTA	AAAAGTACTA	CGGATTATAT
130	140	150	160	170	180
ATCAGTGGAC	AATATAAAC	CAGTGTTC	GTTCAGTA	ATTTTCAGT	TAAAGAAACC
190	200	210	220	230	240
AATGTCATAA	CTAAAAACCT	TATAGCTTTA	AAAAAAGATG	TTGACTCTAT	TGAAACCAAG
250	260	270	280	290	300
ACTGATGCCA	GTGTAGGTAT	TAGTAACCCA	TCAAATTTA	CTATCCCCTA	TACAGCTGTA
310	320	330	340	350	360
TTTCAAGATA	ATTCTGTCAA	TTTCAATGGA	ACTATTGGTT	ACACCTTC	TGAAGGTACA
370	380	390	400	410	420
AGAGTTGAAA	TAGAAGGTT	TTATGAGGAA	TTTGATGTTA	AAAACCTGG	AGGCTATACA
430	440	450	460	470	480
CTAAGTGATG	CCTATCGCTA	TTTGCATTA	GCACGTGAAA	TGAAAGGTA	TAGTTTACA
490	500	510	520	530	540
CCTAAAGAAA	AAGTTCTAA	TAGTATTTT	CACACTGAA	TGAGAAATGA	TGGATTATCT
550	560	570	580	590	600
ATAATATCTG	TTATAGTAAA	TGTTTGCTAC	GATTCTCTT	TGAACAATT	GTCAATATCG
610	620	630	640	650	660
CCTTACATAT	GTGGAGGAGC	AGGGGTAGAT	GCTATAGAAT	TCTTCGATGT	ATTACACATT
670	680	690	700	710	720
AAGTTGCA	ATCAAAGCAA	GCTAGGTATT	GCTTATTCTC	TACCATCTAA	CATTAGCTC
730	740	750	760	770	780
TTTGCTAGTT	TATATTACCA	TAAAGTAATG	GGCAATCAAT	TTAAAATTT	AAATGTCCAA
790	800	810	820	830	840
CATGTTGCTG	AACTTGCAAG	TATACCTAAA	ATTACATCCG	CAGTTGCTAC	ACTTAATATT
850	860	870	880	890	900
GGTTATTTTG GAGGTGAAAT TGGTGCAAGA TTGACATTTT AA.....					

Fig. 25A

10	20	30	40	50	60
MNNKLKFTII	NTVLVCLLSL	PNISSSKAIN	NNAKKYYGLY	ISGQYKPSVS	VFSNFSVKET
70	80	90	100	110	120
NVITKNLIAL	KKDVDSEIETK	TDASVGISNP	SNFTIPTYAV	FQDNSVNENG	TIGYTFAEGT
130	140	150	160	170	180
RVEIEGSYEE	FDVKNPGGYT	LSDAYRYFAL	AREMKGNSFT	PKEKVSNSIE	HTVMRNDGLS
190	200	210	220	230	240
IISVIVNVCY	DFSLNNLIS	PYICGGAGVD	AIEFFDVLHI	KFAYQSKLGI	AYSILPSNISL
250	260	270	280	290	300
FASLYYHKVM GNQFRNLNVQ HVAAELASIPK ITSAVATLNI GYFGGEIGAR LTF.....					

Fig. 25B

10	20	30	40	50	60
ATGGCAAATT	TTATGTACAA	AAAATACAAA	CTAATGACAG	CAGGTGTAGT	ATTATTCAC
70	80	90	100	110	120
ATGTTATTC	TACCTCATGT	TTCTTTCGCA	AAAATACAA	ACAGCAATAA	ACTTGGATTAA
130	140	150	160	170	180
TACATCAGTG	GACAGTATAA	CCCTAGTGT	TCTGTTTTA	GCAATTTTC	AGCAAAAGAA
190	200	210	220	230	240
ACCAATGTT	ATACAGTACA	ACTCATGGCG	CTTAAAAAAG	ACATTGATTC	TATTGAAGTT
250	260	270	280	290	300
GATACTGGAA	ATAGCGCAGG	TATAGCAAA	CCACAAAATT	TCACAGTTCT	TTATACTCCA
310	320	330	340	350	360
AAATTTCAAG	ATAATGTTGC	TGGTCTTAGC	GGTGCACTTG	GATTCTTTA	TTCTAAAGGA
370	380	390	400	410	420
TTAAGGATTG	AAATGGGTT	TTCTTATGAA	AAATTTGATG	CTAAAGACCT	TGGTGAGTAC
430	440	450	460	470	480
ACCAAAATAA	AAGATGCTTA	TAGATATTTT	GCTCTAGTAC	GTGAAATGCA	TGTTAGTCTC
490	500	510	520	530	540
ATTTATCCAA	AAGATAATAA	CACAGGAACA	CATTATACTG	TTATGAGAAA	TGATGGTATA
550	560	570	580	590	600
TCTATTTCTT	CTGCTACAGT	AAATGGCTGC	TATGATTCTT	TTTCCAGTT	TATCTTTGTC
610	620	630	640	650	660
ACCTATATGT	GTATAGGCAT	CGGTATAGAT	GCTATAGAAT	TTCTTAATGC	ATACATATTA
670	680	690	700	710	720
AGTTTGCTTG	CCAAGGTAGT	TAAGGTGTTA	ACTTATTCTG	TATCTCCCAA	TGTTAATTTA
730	740	750	760	770	780
TTTGCAGATG	GATATTATCA	TAAAGTGATG	GGCAATAAAT	TTAAAAAATT	ACCTGTTCAA
790	800	810	820	830	840
TACGTTAATA	CTTTAGAAGA	GTATCCAAGA	GTTACATCTG	CAATTGCTAC	ACTTGATATT
850	860	870	880	890	900
GGCTACCTCG	GTGGTGAAAT	TGGCATAAGA	TTTATATTT	AA.....

Fig. 26A

10	20	30	40	50	60
MYKKYKLMTA	GVVLFHLFL	PHVSFAKNTN	SNKLGLYISG	QYNPSVSVFS	NFSAKETNVH
70	80	90	100	110	120
TVQLMALKKD	IDSIEVDTGN	SAGISKPQNF	TVLYTPKFQD	NVAGLSGALG	FFYSKGLRIE
130	140	150	160	170	180
MGFSYEKFDA	KDLGEYTKIK	DAYRYFALVR	EMHVSЛИYPK	DNNTGTHYTV	MRNDGISIS
190	200	210	220	230	240
ATVNGCYDSE	FQFIEVTVYMC	IGIGIDAIEF	LNAYILSLLA	KVVKVLTYSV	SPNVNLFADG
250	260	270	280	290	300
YYHKVMGNKF	KNLPVQYVNT	LEEYPRVTS	IATLDIGYLG	GEIGIRFIE

Fig. 26B

10	20	30	40	50	60
ATGGGAAATT	CTATGATAA	AAAAAGTC	TTCTTAATAA	GATTTATATT	TTAACATGC
70	80	90	100	110	120
ATGCTGTCAT	TACCTAATAT	ATCTCTTC	AAAGTAAATA	ACGAAAAACA	TTCTGGTTG
130	140	150	160	170	180
TATATTAGCG	GGCAATACAA	ACCCAGTGT	TCTGTTTC	GTAATTTTC	AGTTAAAGAA
190	200	210	220	230	240
ACCAACTTTC	ATACAAAACA	TCTCATAGCT	CTTAAACAAG	ATGTTGATTC	TGTTGAAATT
250	260	270	280	290	300
GATACTGGTA	GTAATACAGC	AGGTATTAGT	ACCCATCTA	ACTTTACAAT	CCCTTATACT
310	320	330	340	350	360
GCAGAAATTC	AAGACAACCA	TACTAACTGC	AATGGCTCTA	TTGGTTATGC	TTTGCTGAA
370	380	390	400	410	420
GGTCCAAGAA	TTGAAATAGA	ATTATCATAT	GAAAAATTG	ATGTTAAAAA	TCCCACAGGG
430	440	450	460	470	480
TATACTACAG	AAAAAGATGC	TTATAGATAC	TTTGCTTAG	CACGTGAAAT	AAATATTTCT
490	500	510	520	530	540
CTATTCCAAC	CAAAACAAAA	AGAAGGTAGT	GGAATTAC	ATGTCGTAAT	GAAAAACGAT
550	560	570	580	590	600
GGGTTATCTA	TCTTATCCAA	TATAGTTAAT	ATTTGCTACG	ATTTTCTTT	AAATAATTAA
610	620	630	640	650	660
CCTATATCAC	CTTATTTATG	CGGAGGAATG	GGTATAAAATG	CCATAGAATT	CTTGACGCT
670	680	690	700	710	720
TTACATGTGA	AATTTGCTTA	TCAAAGCAAG	GCAGGAATTA	GTTATCAACT	ATTACGTAAA
730	740	750	760	770	780
ATCAACTTAT	TTATTGATGT	ATATTACTAC	GAAGTAATAA	GTAATAAATT	AAAAAACCTG
790	800	810	820	830	840
AAAGTCCAAC	ATGTACATGA	ACTTAAAGAT	AATCCAAAAG	TCACATCTGC	AGTTGCTACA
850	860	870	880	890	900
CTTGATATAG CATATTTGG TAGTGAAGCT GGCATAAGAA TTATATTTA A.....					

Fig. 27A

10	20	30	40	50	60
MNNKSQFLIR	FIFLTCLMSL	PNISLSKVNN	EKHSGLYISG	QYKPSVSVFS	NFSVRETNFH
70	80	90	100	110	120
TKHLIALKQD	VDSVEIDTGS	NTAGISNPSN	FTIPTYAEOFQ	DNHTNCNGSI	GYAFAEGPRI
130	140	150	160	170	180
EIELSYEKFD	VKNPTGYTTV	KDAYRYFALA	REINISLFQP	KQKEGSGIYH	VVMKNDGLSI
190	200	210	220	230	240
LSNIVNICYD	FSLNNLPISP	YLCGGMGINA	IEFFDALHVK	FAYQSKAGIS	YQLLRKINLE
250	260	270	280	290	300
IDVYYEVIS NKFKNLKVQH VHELKDNPKV TSAVATLDIA YFGSEAGIRI IF.....					

Fig. 27B

10	20	30	40	50	60
ATGAATAGCA	AGAGTAAGTT	CTTTACAATA	TGTACATCGT	TAATATGCCT	ATTATCATCA
70	80	90	100	110	120
CCTAACACAT	CTCTCTCAAA	CTTCATAGGC	AATAGTACAA	AACATTCTGG	ATTATATGTT
130	140	150	160	170	180
AGCGGACAAT	ATAAGCCCAG	CGTTTCCATT	TTTAGCAAAT	TTTCAGTAAA	AGAAACAAAT
190	200	210	220	230	240
ACACATACAG	TACAGTTAGT	AGCTCTTAAA	AAAGATGTTA	ATTCTATTTC	TATGAACATC
250	260	270	280	290	300
AGTAATGGTG	CTACAGGCAT	TAGCAAAGCA	ACAAATTTA	ATCTTCCTTA	TGTTGCAGAA
310	320	330	340	350	360
TTTCAAGACA	ATGCCTTCAA	CTTCAGTGGG	GCTATTGGTT	ATTCACTTTT	TGAACAACTA
370	380	390	400	410	420
AACATTGAAG	TGGAAGGTTG	TTATGAAGAA	TTCGATGCCA	AAAATCCTGG	TGGTTATATT
430	440	450	460	470	480
TTAAATGATG	CATTCCGCTA	TTTTGCATTG	GCACGTGAAA	TGGGACAAGA	AAAAAATGAT
490	500	510	520	530	540
AATAAGCATC	TTAGTCCTAA	GGAGGAGCAT	GATATAAGTA	AAACATATTAA	CACAGTCATG
550	560	570	580	590	600
AGAAAATAATG	GGTTATCTAT	ATTATCTATT	ATGATAAAATG	GCTGCTATAA	TCTACCTCTC
610	620	630	640	650	660
AATGATTTAT	CAATATCACC	TTATTTTGT	ACAGGAATAG	GTGTAGATGC	TATAGAATTT
670	680	690	700	710	720
TTTGATGCAC	TGCATCTTAA	ACTTGCTTG	CAAAAGTAAAA	TAGGAGCTAC	TTACCAATTAA
730	740	750	760	770	780
TCAGACAAACA	TTAGTTTATT	TACAARTGGA	TATTACCATC	AAGTAATAGG	TGATCAATTT
790	800	810	820	830	840
AAAAAACTTAA	AAGTCCAATA	TATAGGTGAA	CTTAAAGAGA	ACCCGAAAAT	TACATCTGCA
850	860	870	880	890	900
GTTGCTACTC	TCAATGTTGG	ATACTTTGGA	GGTGAATATG	GAGTAAGACT	CACACTTTAA
910	920	930	940	950	960

Fig. 28A

10	20	30	40	50	60
MNSKSKEFTI	CTSLICLLSS	PNTSLSMFIG	NSTKHSGLYV	SGQYKPSVSI	FSKESVKETN
70	80	90	100	110	120
THTVQLVALK	KDVNSISMNI	SNGATGISKA	TNFNLPYVAE	FQDNAFNESG	AIGYSLFEQL
130	140	150	160	170	180
NIEVEGSYEE	FDAKNPGGYI	LNDAFRYFAL	AREMGQEKNND	NKHLSPKEEH	DISKTYYTVM
190	200	210	220	230	240
RNNGLSILSI	MINGCYNLPL	NDLSISPYFC	TGIGVDAIEF	FDALHLKLAL	QSKIGATYQL
250	260	270	280	290	300
SDNISLFTNG	YYHQVIGDQE	KNLKVOYIGE	LKENPKITSA	VATLNVGYFG	GEIGVRLTL.

Fig. 28B

10	20	30	40	50	60
AAGCTTCTTA	TGAAGAATTT	GACGTTAAAA	ATCCTGAAGG	ATCTACTACA	GAACCTATA
70	80	90	100	110	120
GATATTTCGC	GTTAGCACGT	GGCATGGATG	GTAATAATAT	TCCTACAAGT	CAAAAATTTA
130	140	150	160	170	180
CTGTAATGAG	AAACGACGGG	TTATTAATCT	CATCTGTTAT	GATAAATGGC	TGTTACAATG
190	200	210	220	230	240
TCATACTAAA	TGATATACAA	GCAGAACCTT	ACATATGTGC	AGGACTAGGA	GGAGATTTA
250	260	270	280	290	300
TAGAATTCTT	CAATGGCTTT	CATGTTAACG	TAGCTTATCA	AGGTAAAGTA	GGCATTAGTT
310	320	330	340	350	360
ATCAAATATT	CCCTGAAGTA	AGATTATTAA	TTGATGGATA	CTACCATAAA	GTAAAAGGCA
370	380	390	400	410	420
ACAAGTTAA	AAATTTACAC	GTTCAACATG	TAGGTGCACT	TGCAGCACTC	CCTAAAGTTA
430	440	450	460	470	480
CATCTGCAGT	TGCAACACTT	AATATTGGAT	ACTTTGGTTG	TGAAGCTGGA	GTAAGATTCA
490	500	510	520	530	540
TATTTTAA..

Fig. 29A

10	20	30	40	50	60
ASYEEFDVKN	PEGSTTDSYR	YFALARGMDG	NNIPTSQKFT	VMRNDGLLIS	SVMINGCYNV
70	80	90	100	110	120
ILNDIQAEPY	ICAGLGGDFI	EFFNGFHVKL	AYQGKVGISY	QIFPEVRLFI	DGYYHKVKGN
130	140	150	160	170	180
KFKNLHVQHV	GALAALPKVT	SAVATLNIGY	FGCEAGVRFI	F.....

Fig. 29B

29/31

10	20	30	40	50	60
ATGAAATTATA	AGAAAATTCT	AGTAAGAAGC	GCGTTAACCT	CATTAATGTC	AATCTTACCA
70	80	90	100	110	120
TATCAGTCCTT	TTGCAGATCC	TGTAGGTTCA	AGAACTAACATG	ATAACAAAGA	AGGCTTCTAC
130	140	150	160	170	180
ATTAGTGCAA	AGTACAATCC	AAGTATATCA	CACTTTAGAA	AATTCTCTGC	TGAAGAAACT
190	200	210	220	230	240
CCTATTAATG	GAACAAATTC	TCTCACTAAA	AAAGTTTCG	GACTAAAGAA	AGATGGTGAT
250	260	270	280	290	300
ATAACAAAAA	AAGACGATTT	TACAAGAGTA	GCTCCAGGCA	TTGATTTCA	AAATAACTTA
310	320	330	340	350	360
ATATCAGGAT	TTTCAGGAAG	TATTGGTTAC	TCTATGGACG	GACCAAGAAT	AGAACTTGAA
370	380	390	400	410	420
GCTGCATATC	ACAATTTAAT	CCAAAAACAC	GATAACAATG	ATACTGATAA	TGGTGAATAC
430	440	450	460	470	480
TATAAACATT	TTGCATATCT	CGTAAAGATG	CCATGGAAGA	TCAGCCATAT	GTTGTTCTTA
490	500	510	520	530	540
AAAATGACGG CATA.....					

Fig. 30A

10	20	30	40	50	60
MNYKKILVRS	ALISLMSILP	YQSFADPVGS	RTNDNKEGFY	ISAKYNPSIS	HERKFSAEET
70	80	90	100	110	120
PINGTNSLT	KVFGLEKKDGD	ITKKDDFTRV	APGIDFQNNL	ISGFSGSIYG	SMDGPRIELE
130	140	150	160	170	180
AAYHNLIQKH DNNNDTNGEY YKHFAYLVKM PWKISHMLFL KMTAY.....					

Fig. 30B

30/31

SV												
CNP-17												
CNP-18												
CNP-19												
CNP-1C												
CNP-1B												
P20												
HAP-1												
CNP-1A												
MV1												
CNP-17												
CNP-18												
CNP-19												
CNP-1C												
CNP-1B												
P20												
HAP-1												
CNP-1A												
MV2												
CNP-17												
CNP-18												
CNP-19												
CNP-1C												
CNP-1B												
P20												
HAP-1												
CNP-1A												
MV3												
CNP-17												
CNP-18												
CNP-19												
CNP-1C												
CNP-1B												
P20												
HAP-1												
CNP-1A												

Fig. 31

31/31